



Cover: Primary cortical neuronal culture isolated from newborn Tg2576 mice, expressing the human amyloid precursor protein (APP, isoform 695) carrying the Swedish mutation (APP^{Sw}; KM670/671NL). Cells were allowed to mature *in vitro* for 8 days and then characterized for culture composition by immunocytochemistry for neural (β -III-tubulin; green) and astroglial (GFAP; red) markers. See article by Baldassarre et al. on page 671. Cover image by Vito Antonio Baldassarre is licensed under a Creative Commons Attribution 4.0 International licence.

EDITORIAL

- 499** Neurodegenerative disease: models, mechanisms, and a new hope
Gitler, A. D., Dhillon, P. and Shorter, J.

A MODEL FOR LIFE

- 503** Solving the puzzle of neurological diseases: an interview with Huda Zoghbi
Zoghbi, H. Y.

AT A GLANCE

- 509** RNA metabolism in neurodegenerative disease
Liu, E. Y., Cali, C. P. and Lee, E. B.

REVIEWS

- 519** The role of Ca^{2+} signaling in Parkinson's disease
Zaichick, S. V., McGrath, K. M. and Caraveo, G.

- 537** Modelling amyotrophic lateral sclerosis: progress and possibilities
Van Damme, P., Robberecht, W. and Van Den Bosch, L.

RESEARCH ARTICLES

- 551** Upregulation of CB₂ receptors in reactive astrocytes in canine degenerative myelopathy, a disease model of amyotrophic lateral sclerosis
Fernández-Trápero, M., Espejo-Porras, F., Rodríguez-Cueto, C., Coates, J. R., Pérez-Díaz, C., de Lago, E. and Fernández-Ruiz, J.

- 559** Loss of Ranbp2 in motoneurons causes disruption of nucleocytoplasmic and chemokine signaling, proteostasis of hnRNPH3 and Mmp28, and development of amyotrophic lateral sclerosis-like syndromes
Cho, K.-i., Yoon, D., Qiu, S., Danziger, Z., Grill, W. M., Wetsel, W. C. and Ferreira, P. A.

- 581** The tyrosine kinase receptor Tyro3 enhances lifespan and neuropeptide Y (Npy) neuron survival in the mouse anorexia (*anx*) mutation
Kim, D. Y., Yu, J., Mui, R. K., Niibori, R., Taufique, H. B., Aslam, R., Semple, J. W. and Cordes, S. P.

597

Schwann cells are activated by ATP released from neurons in an *in vitro* cellular model of Miller Fisher syndrome

Rodella, U., Negro, S., Scorzetto, M., Bergamin, E., Jalink, K., Montecucco, C., Yuki, N. and Rigoni, M.

605

The familial dysautonomia disease gene *IKBKP* is required in the developing and adult mouse central nervous system

Chaverra, M., George, L., Mergy, M., Waller, H., Kujawa, K., Murnion, C., Sharples, E., Thorne, J., Podgajny, N., Grindeland, A., Ueki, Y., Eiger, S., Cusick, C., Babcock, A. M., Carlson, G. A. and Lefcort, F.

619

Blood RNA biomarkers in prodromal PARK4 and rapid eye movement sleep behavior disorder show role of complexin 1 loss for risk of Parkinson's disease

Lahut, S., Gispert, S., Ömür, Ö., Depboylu, C., Seidel, K., Domínguez-Bautista, J. A., Brehm, N., Tireli, H., Hackmann, K., Pirkevi, C., Leube, B., Ries, V., Reim, K., Brose, N., den Dunnen, W. F., Johnson, M., Wolf, Z., Schindewolf, M., Schrempf, W., Reetz, K., Young, P., Vadasz, D., Frangakis, A. S., Schröck, E., Steinmetz, H., Jendrach, M., Rüb, U., Başak, A. N., Oertel, W. and Auburger, G.

633

Genetic mutations linked to Parkinson's disease differentially control nucleolar activity in pre-symptomatic mouse models

Evsyukov, V., Domanskyi, A., Bierhoff, H., Gispert, S., Mustafa, R., Schlaudraff, F., Liss, B. and Parlato, R.

645

A differential autophagy-dependent response to DNA double-strand breaks in bone marrow mesenchymal stem cells from sporadic ALS patients

Wald-Altmann, S., Pichinuk, E., Kakhlon, O. and Weil, M.

655

Cellular levels of Grb2 and cytoskeleton stability are correlated in a neurodegenerative scenario

Majumder, P., Roy, K., Singh, B. K., Jana, N. R. and Mukhopadhyay, D.

671

Vulnerability of primary neurons derived from Tg2576 Alzheimer mice to oxygen and glucose deprivation: role of intraneuronal amyloid- β accumulation and astrocytes

Baldassarre, V. A., Marchesini, A., Giardino, L. and Calzà, L.