

REVIEW

Modeling congenital kidney diseases in *Xenopus laevis*

Alexandria T. M. Blackburn^{1,2} and Rachel K. Miller^{1,2,3,4,*}

ABSTRACT

Congenital anomalies of the kidney and urinary tract (CAKUT) occur in ~1/500 live births and are a leading cause of pediatric kidney failure. With an average wait time of 3-5 years for a kidney transplant, the need is high for the development of new strategies aimed at reducing the incidence of CAKUT and preserving renal function. Next-generation sequencing has uncovered a significant number of putative causal genes, but a simple and efficient model system to examine the function of CAKUT genes is needed. *Xenopus laevis* (frog) embryos are well-suited to model congenital kidney diseases and to explore the mechanisms that cause these developmental defects. *Xenopus* has many advantages for studying the kidney: the embryos develop externally and are easily manipulated with microinjections, they have a functional kidney in ~2 days, and 79% of identified human disease genes have a verified ortholog in *Xenopus*. This facilitates high-throughput screening of candidate CAKUT-causing genes. In this Review, we present the similarities between *Xenopus* and mammalian kidneys, highlight studies of CAKUT-causing genes in *Xenopus* and describe how common kidney diseases have been modeled successfully in this model organism. Additionally, we discuss several molecular pathways associated with kidney disease that have been studied in *Xenopus* and demonstrate why it is a useful model for studying human kidney diseases.

KEY WORDS: *Xenopus*, Nephron, CAKUT, Nephronophthisis, PKD, CRISPR, Kidney

Modeling CAKUT in *Xenopus*

Congenital anomalies of the kidney and urinary tract (CAKUT; see Box 1 for a glossary of terms) are a leading cause of pediatric kidney failure, accounting for 40-50% of pediatric chronic kidney disease (CKD; Box 1) worldwide (Vivante et al., 2014). CAKUT encompasses a wide range of structural malformations resulting from morphogenetic defects, including Wilms tumor and renal hypodysplasia (Box 1; Fig. 1) (Yosypiv, 2012). With an average 3- to 5-year wait time for a deceased donor kidney transplant [United Network for Organ Sharing (<https://www.kidney.org/atoz/content/transplant-waitlist>), accessed December 10, 2018], the need to find alternative treatments that preserve renal function is essential. Monogenic disease with strong genetic causality only accounts for

12% of CAKUT cases (Vivante and Hildebrandt, 2016), and polygenic causes are speculated to occur but are largely unknown (Sanna-Cherchi et al., 2018). Next-generation sequencing has helped to uncover novel causative genes of CAKUT, but a high-throughput strategy to test their function in the kidney is needed. To understand how CAKUT arises, it is crucial to understand how the kidneys and urinary tract develop to uncover the genetic mechanisms that coordinate these events. Mice and zebrafish have been the predominant models used in kidney research, while recent advances have made kidney organoids (Box 1) useful for nephrotoxicity screening as well as for modeling kidney diseases (Hurtado del Pozo et al., 2018). However, *Xenopus laevis* frogs (Box 1; hereafter referred to simply as *Xenopus*) possess many qualities that make them an effective *in vivo* model to study congenital kidney diseases.

Xenopus share a relatively close evolutionary history with mammals because frogs are tetrapods. Thus, this model has the advantage of rapid development, like zebrafish, but evolutionarily it lies closer to mammals. Additionally, *Xenopus* and human genomes have long stretches of gene collinearity, and 79% of identified human disease genes have a verified ortholog in *Xenopus* (Hellsten et al., 2010). The embryonic kidney of *Xenopus* has many characteristic features of a mature mammalian kidney (Fig. 2) (Zhou and Vize, 2004; Raciti et al., 2008). Thus, many genes and processes necessary for *Xenopus* kidney development are also important in mammalian kidney development. Furthermore, a limited number of kidney-disease-causing genes have been analyzed in *Xenopus* or are expressed in its embryonic kidney, introducing potential paths for future kidney research (Fig. 1 and Table 1). Although few in number, the *Xenopus* genes studied have been shown to function in kidney development, as their mammalian orthologs do in mammalian kidneys, and their disrupted expression results in similar nephron (Box 1) phenotypes to those seen in humans. Additionally, *Xenopus* possesses unique qualities that other *in vivo* model systems lack.

With a simple hormone injection, *Xenopus* produces large clutch sizes with hundreds of embryos that develop externally. Their kidneys can be easily visualized and imaged through their transparent epidermis, and they develop a fully functional kidney in ~56 hours post-fertilization (hpf) (Vize et al., 2003). Additionally, cleavage-stage *Xenopus* embryos have been fate-mapped (Box 1), which allows for the tracking of specific blastomeres (Box 1) as they develop from an early-stage embryo into a differentiated body plan (Dale and Slack, 1987; Moody, 1987). This facilitates unilateral tissue-targeted injections that are specific to organs such as the kidney (DeLay et al., 2016) and permits using the uninjected side as an internal control. In addition, researchers have developed assays to study kidney function in *Xenopus* (Zhou and Vize, 2004; Tran et al., 2010).

Xenopus oocytes (Box 1) and early embryos can easily be injected with DNA, mRNA, protein and/or morpholinos (Box 1) to overexpress or knock down proteins. A common assay in *Xenopus* involves disrupting genes/proteins in both kidneys by injecting

¹Pediatric Research Center, Department of Pediatrics, McGovern Medical School, The University of Texas Health Science Center, Houston, TX 77030, USA. ²The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences, Program in Genetics and Epigenetics, Houston, TX 77030, USA. ³The University of Texas MD Anderson Cancer Center UTHealth Graduate School of Biomedical Sciences, Program in Biochemistry and Cell Biology Houston, Houston, TX 77030, USA. ⁴Department of Genetics, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA.

*Author for correspondence (Rachel.K.Miller@uth.tmc.edu)

© A.T.M.B., 0000-0003-2974-0339; R.K.M., 0000-0002-4214-3347

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium provided that the original work is properly attributed.

Box 1. Glossary

Ciliopathy: a genetic disorder caused by abnormal formation or function of cilia (a component of almost all cells). Disruption of cilia leads to a recognizable list of features, including retinal degeneration, cardiac defects, mental retardation and kidney disease.

Epigenetic signature: a set of epigenetic marks, such as methylation, found on specific genes that are associated with the phenomenon being observed, such as a disease state.

Excretion assay: fluorescent dyes, such as rhodamine, can be injected into the *Xenopus* coelomic cavity and filtered by the kidneys. These dyes are then secreted into the urine, which allows for a simple visual readout of kidney function.

Expressivity: measures the extent to which a given genotype manifests (is expressed) at the phenotypic level. It accounts for different degrees of phenotypic expression in different individuals, which may be due to environmental factors or the allelic constitution of the rest of the genome.

Fate mapping: determines what types of cells, tissues and organs are derived from specific embryonic cells. Classical fate maps inject a lineage tracer such as a fluorescent dextran into a specific cell, which then allows all of its descendants to retain the fluorescence and therefore be mapped.

GAL4-UAS: involves the development of two lines: the GAL4 line, which expresses the transcription factor GAL4 in a subset of the animal's tissues; and the UAS line, in which UAS is normally expressed upstream of fluorescent proteins and acts as a reporter. GAL4 specifically binds to UAS promoter elements, thus activating expression of the downstream target sequence.

Genome-wide association study (GWAS): an observational study of the entire genome and its set of genetic variants in different individuals to see whether any variant may be associated with a specific trait. These studies generally focus on single-nucleotide variants and associations that may lead to a predisposition of various diseases.

Glomerulopathy: a disease affecting the glomeruli of the nephron that causes the kidneys to malfunction. Features include high levels of protein and sometimes blood in the urine, and swelling in many areas of the body. Loss of glomerular filtration leads to end-stage renal disease (ESRD) in about half of the individuals within 10 years of their diagnosis.

Heat-shock inducible: a heat shock promoter, normally the *Hsp70* promoter, is used to regulate transgene expression when the ambient temperature is briefly increased. This heat shock releases a factor that then allows it to bind to elements of the promoter, thus activating transcription. This technique has been used in aquatic animals such as zebrafish and *Xenopus*, as well as in *Caenorhabditis elegans* and *Drosophila*.

Intermediate mesoderm: part of the mesoderm germ layer that is located between the paraxial and lateral plate mesoderm. The intermediate mesoderm gives rise to the reproductive and urogenital systems, including the kidney.

Kidney organoids: result from kidney organogenesis in a dish (*in vitro*) by inducing human pluripotent stem cells to a kidney fate. The organoids contain cell clusters that express markers of various regions of the kidney, such as podocytes, proximal tubules, loops of Henle and the distal tubule. Additionally, *Xenopus* explants of early embryos can be induced to form kidney organoids in culture.

Leapfrogging: transplanting the germline of a *Xenopus* embryo that has been mutagenized (such as with CRISPR/Cas9) into a wild-type host that had its wild-type germline removed. This results in the efficient transmission of mutant alleles to F₁ offspring and overcomes the embryonic lethality of various gene knockouts in the F₀ embryos.

Meckel-Gruber syndrome: a rare autosomal recessive ciliopathy characterized by renal cystic dysplasia, polydactyly and central nervous system malformations. Most individuals with Meckel-Gruber syndrome die before or shortly after birth.

Mesonephros: Greek for 'middle kidney'; the main excretory organ of aquatic vertebrates and a temporary kidney in reptiles, birds and mammals. It develops posteriorly and replaces the pronephros. In humans, the mesonephros functions between the sixth and tenth weeks of embryological life.

Metanephric mesenchyme (MM): all of the cells present in mature nephrons arise from the MM. During kidney development, cells of the MM condense around the ureteric bud to form what's known as the cap

mesenchyme, which goes on to form the renal vesicles, then the comma-shaped bodies and so forth.

Metanephros: the third stage of kidney development, which corresponds to the mature and functional kidney in reptiles, birds and mammals. In humans, the metanephros develops by the tenth week of embryological life, replacing the mesonephros.

Morpholino: antisense oligonucleotide that binds to complementary RNA to knock down gene expression. Its molecular structure contains DNA bases that are attached to a backbone of methylmorpholine rings linked to non-ionic phosphorodiamidate linkages instead of anionic phosphodiester ones, which make them highly stable.

Nail-patella syndrome: an autosomal-dominant disease marked by poorly developed nails, kneecaps (patellae), elbows and pelvis, as well as kidney disease. Nail-patella syndrome is caused by mutations in *LMX1B*.

Nephric primordia: the primordia is the simplest set of cells that are specified to become the kidney and the earliest recognizable stage of kidney development; also known as pronephric anlage.

Nephron: the structural and functional unit of the kidney that filters body waste and excess fluid. The nephron is composed of a glomerulus to filter blood; proximal tubules, a loop of Henle and distal tubules for reabsorbing water and various nutrients; and a collecting duct that resorbs water and carries urine from the kidneys to the bladder.

Nephronophthisis (NPHP): an autosomal-recessive ciliopathy that generally occurs as an isolated kidney disease, although ~15% of NPHP patients also present with extrarenal symptoms. Features include small cysts in the kidney medulla and kidney fibrosis, with three clinical forms classified by the onset of ESRD: infantile, juvenile and adolescent.

Nephrostomes: multi-ciliated cells that funnel blood filtrate from the coelomic cavity and into the tubules of the *Xenopus* kidney.

Neurula: the embryonic stage of development in which neurulation, the transformation of the neural plate into the neural tube, occurs. This process begins when the notochord induces the formation of the central nervous system by forming the neural plate.

Oocyte: an immature female egg that matures within a follicle in the outer layer of the ovaries. *Xenopus* oocytes have been commonly used to study ion transport and channel physiology because of their large diameter (1 mm), as well as making cell-free extracts to study cell and molecular biology.

Polycystin signaling: biochemical interactions between proteins that lead to cyst formation in the kidney, including HNF1B, TSC2 and BICC1, and the *PKD1* and *PKD2* gene products, polycystin 1 and polycystin 2. Cyst formation may occur due to their downstream signaling pathways or due to the direct interactions between these proteins.

Pronephros: the first stage of kidney development that is functional in aquatic vertebrates such as *Xenopus*, but nonfunctional in reptiles, birds and mammals. In humans, the pronephros is a vestigial structure that disappears completely by the fourth week of embryonic life. Despite this transient appearance in mammals, the pronephros is essential for the development of the adult kidneys.

Renal vesicles: the primordial structure of the nephron that is the first polarized epithelial derivative of the metanephric mesenchyme. Renal vesicles sequentially evolve into the comma-shaped body, followed by the S-shaped body and finally the nephron.

Retinoic acid signaling: a concentration-dependent signaling pathway important for embryo patterning and development. Retinoic acid signaling is essential for kidney development and regulates embryonic kidney patterning.

Septins: a group of GTP-binding proteins that form complexes including filaments and rings. Septins are a unique cytoskeletal component that have been implicated in the localization of proteins at the cell membrane where cilia are found.

S-shaped bodies: the epithelial precursors for the nephron segments (glomerulus, proximal tubule and distal tubule). The segments are oriented along the proximal-distal axis with the glomerulus being the most proximal.

Tet-On: an inducible gene expression system that requires two lines, one carrying the gene of interest under the control of a tetracycline-inducible promoter (fusion of CMV minimal promoter and seven copies of the *tetO* sequence) and the other carrying the transcriptional activator, rTA. In the

Continued on next page

Box 1. Glossary (continued)

absence of rTA, the CMV minimal promoter cannot drive the expression of the transgene. However, when rTA binds to the *tetO* sequences, RNA polymerase can be recruited to the CMV minimum promoter to begin transcription.

Tubulopathy: a disease that affects the tubules of the nephrons in the kidney. It can arise from mutations in ion channel genes such as *CLCNKB*. Tubular dysfunction can cause profound electrolyte and volume disturbance.

Wilms tumor: a type of childhood cancer in the kidneys that most often affects children aged 3 to 4 years. Most Wilms tumors only affect one kidney and the chance of curing children with these tumors is very high.

Wnt pathway: a group of signaling pathways that are essential for embryonic development and are highly conserved in animals, from fruit

flies to humans. There are three main branches of the Wnt pathway: the canonical Wnt pathway involving β -catenin, the non-canonical planar cell polarity pathway, and the non-canonical Wnt/calcium pathway. All three pathways are activated when a Wnt ligand binds to a Frizzled receptor and transduces the signal to the Dishevelled protein. Both canonical and non-canonical planar cell polarity pathways are important for kidney development, with potential involvement of the calcium pathway (Halt and Vainio, 2014).

***Xenopus laevis*:** an aquatic species of African clawed frogs that have been used as a model organism to study embryonic development, cell and molecular biology, as well as in large-scale genetic screens. *Xenopus laevis* is an established model of nephron development.

morpholinos, or CRISPR sgRNAs and the Cas9 protein, in both ventral cells of four-cell embryos. If the gene/protein is important for kidney formation, its loss will result in edema, characterized by swelling in the chest cavity due to fluid retention. This technique allows for disruption of the kidney while avoiding the heart and liver, two other common causes of edema, generating tissue-targeted knockdown or knockout embryos. Upon edema formation, researchers can then use an excretion assay (Box 1) that allows

visualization of the passage of fluorescent molecules through the kidney and out through the cloaca, which can show whether the kidneys are still functioning (Zhou and Vize, 2004). Although not directly comparable to the *Cre-loxP* systems used in mouse studies, this edema assay, which is unique to *Xenopus*, allows for some evaluation of tissue specificity. Knockdown of *Pkd2*, a protein involved in autosomal-dominant polycystic kidney disease (ADPKD; Box 1), in *Xenopus* embryonic kidneys results in




Fig. 1. Common malformations of the kidney found in CAKUT. Renal malformations resulting from inherited kidney diseases are depicted in colored boxes. The colors correspond to the color coding for genes whose loss results in the given phenotype. (1) Green represents renal cysts that are large and cover the majority of the kidney, as seen in renal dysplasia, multicystic dysplastic kidney (MCDK) and autosomal-dominant polycystic kidney disease (ADPKD). (2) Magenta represents a tumor, as seen in tuberous sclerosis and Wilms tumor. (3) Purple represents kidney agenesis (Box 1). (4) Teal represents renal hypoplasia, which is one of the most common CAKUT phenotypes. (5) Orange represents nephronophthisis, with maroon spots depicting corticomedullary cysts, which are generally small. (6) Blue represents the horseshoe kidney, where both kidneys are fused together. (7) Red represents ureter malformations and blockages, which result in urine backflow into the kidney (shown in yellow in the schematic). Genes listed in the key have been studied or are expressed in the *Xenopus* kidney. Numbers found in this figure in the bottom left corner that correspond to the aforementioned phenotypes can also be found in Table 1 under 'Renal phenotype'.




Fig. 2. *Xenopus* pronephric and human metanephric nephrons share conserved tubule segmentation patterns based on gene expression data.

The schematic represents nephron segments, designated by different colors, in the mammalian metanephric nephron (top) and the *Xenopus* pronephric nephron (bottom). The glomus/glomerulus filters blood across capillary walls into the proximal tubule, which filters various wastes out of the body through the remaining distal and connecting tubules. There are noted differences between the glomus and glomerulus in that the glomus deposits blood filtrate into the coelomic cavity. Additionally, *Xenopus* does not have a loop of Henle (grayed out in human schematic) or a true collecting duct (grayed out in human schematic), but instead has a region analogous to the connecting tubule closest to the distal tubule of the mammalian metanephric nephron (Raciti et al., 2008).

edema (Tran et al., 2010). Similar edema phenotypes occur due to the loss of other proteins implicated in CAKUT, such as Pax2 and Pax8 (Buisson et al., 2015). These established techniques demonstrate that *Xenopus* possesses distinctive qualities suitable for disease modeling, and new technologies continue to improve the experimental opportunities in this model system.

Genetic screens of the allotetraploid (Box 1) *Xenopus* are becoming more feasible with the development of the CRISPR/Cas9 system (Aslan et al., 2017), and the ability to target CRISPR-mediated genetic manipulation to the kidney is unique to *Xenopus* (DeLay et al., 2018). These advantages make *Xenopus* a valuable model for studying the kidney, can aid in our understanding of the mechanisms through which the kidney develops, and can help identify new genes important for renal function. This Review will highlight the *Xenopus* genes and pathways that are associated with human kidney disease, demonstrate the similarity in phenotypes upon gene or protein disruption, and establish *Xenopus* as a relevant model for CAKUT.

Kidney development

The kidney is an excretory organ that filters blood. It functions by reabsorbing nutrients and excreting waste and excess fluid in the form of urine. In humans, nephrogenesis starts at around 5 weeks of gestation and completes between weeks 34 and 36 (Black et al., 2012). Environmental factors, such as vitamin A deficiency or exposure to teratogenic substances such as angiotensin-converting enzyme (ACE) inhibitors, among others, can affect proper kidney development (Rosselot et al., 2010; Al-Maawali et al., 2012). Retinoic acid is derived from vitamin A and retinoic acid signaling (Box 1) is important for embryonic kidney patterning (Rosselot et al., 2010). Additionally, taking teratogenic ACE inhibitors during pregnancy can lead to CAKUT, as the kidney also plays an essential role in controlling blood pressure (Al-Maawali et al., 2012). The kidney is a unique organ in that it develops from successive forms

that replace the previous structure (Davidson, 2008). These forms are the pronephros, mesonephros and metanephros (Box 1). The organizational complexity increases as each sequential form of the kidney is superseded by the next (Davidson, 2008). In mammals, both the pro- and mesonephros are replaced by the metanephros, which persists to function as the adult kidney (Little and McMahon, 2012). In contrast, the onset of metamorphosis in *Xenopus* marks the replacement of the pronephros with the mesonephros, which is the adult kidney of amphibians. It is important to note that, although the pronephros is non-functional in mammals, it is required for the subsequent formation of the mesonephros and metanephros.

Although the arrangement of kidneys differ, the structural and functional unit of the kidney, the nephron, remains the same. An adult human kidney contains approximately 1 million nephrons (Bertram et al., 2011), whereas an adult mouse kidney has over 10,000 (Short et al., 2014). Mammalian nephrons develop asynchronously, making them challenging to study (Sorokin et al., 1992). In contrast, *Xenopus* embryos contain only one functional nephron on either side of their body, which serves as the embryonic kidney. Therefore, the fully functional *Xenopus* pronephric nephron serves as a simplified model for mammalian meso- and metanephric nephron development (Fig. 2).

Conservation of the nephron in *Xenopus* and mammals

The pronephros forms from the intermediate mesoderm (Box 1) in both mammals and amphibians. In *Xenopus*, the nephric primordia (Box 1) is induced by signals from the surrounding tissues and forms during neurula (Box 1) stages 12–15 (~16–20 hpf), and later generates the tubules of the pronephros (Fox, 1963). Pronephros specification in *Xenopus* occurs through the interactions between the transcription factors *Osr1*, *Osr2*, *Pax8*, *Lhx1* and *Hnf1B*, among others (Vize et al., 2003; Tena et al., 2007; Drews et al., 2011; Boualia et al., 2013). Morphogenesis of the pronephros begins at the early-tailbud stage 21 (~23 hpf) (<http://www.xenbase.org/anatomy/alldev.do>). In mice, the *Osr1* gene is essential for the formation of renal structures and can be found in the intermediate mesoderm, while *Osr2* is found later on in the mesonephros and is not essential (Tena et al., 2007). Consistent with *Xenopus* (Buisson et al., 2015; DeLay et al., 2018), mice that lack *Lhx1* or both *Pax2* and *Pax8* have severe kidney defects (*Pax2* and *Pax8* seem to have redundant functions in mice) (Shawlot and Behringer, 1995; Bouchard et al., 2002). Finally, *Hnf1B* is expressed in the ureteric bud as well as in comma- and S-shaped bodies (Box 1), and mutations in this gene lead to kidney cyst formation in mice (Gong et al., 2009). In addition, *Wnt* (wingless/integrated; Box 1), *FGF* (fibroblast growth factor), *BMP* (bone morphogenetic protein) and *GDNF* (glial-cell-line-derived neurotrophic factor) signaling pathways are involved in both *Xenopus* and mammalian kidney development (Dressler, 2006; Costantini and Kopan, 2010).

Analogous to the mammalian nephron, the amphibian pronephros is segmented along its proximal-distal axis (Møbjerg et al., 2000). Mammals have a glomerulus to filter blood, proximal tubules, a loop of Henle and distal tubules for reabsorbing water and various nutrients, and a collecting duct that reabsorbs water and carries urine from the kidneys to the bladder (Little and McMahon, 2012). Similarly, *Xenopus* embryos have a glomus for filtering blood, proximal and distal tubules for reabsorbing nutrients and a connecting tubule, which opens out to the cloaca (Fig. 2). In addition, at the tips of the proximal tubules, the amphibian also has multi-ciliated cells called nephrostomes (Box 1), which cause fluid influx from the coelomic cavity (Box 1) into the proximal tubules (Wessely and Tran, 2011).

Table 1. CAKUT genes that have been analyzed or have kidney expression in *Xenopus*

Gene symbol	Renal phenotype	Human disease	Findings in <i>Xenopus</i>	Mode of inheritance	References
<i>ACE</i>	Renal dysgenesis	Renal tubular dysgenesis	ACE activity detected at high levels in kidney and testes	Recessive	Quassinti et al., 2007; Gribouval et al., 2012
<i>AGT</i>	Renal dysgenesis	Renal tubular dysgenesis	Mitochondrial targeting in liver cells	Recessive	Holbrook and Danpure, 2002; Gribouval et al., 2012
<i>ANKS6 (NPHP16)</i>	Renal hypoplasia ⁴ , renal cysts ⁵	Nephronophthisis, situs inversus	Role in early tubule morphogenesis	Recessive	Hoff et al., 2013; Taskiran et al., 2014
<i>BICC1</i>	Renal cysts ¹ , renal dysplasia ¹	Renal disease	Regulates renal epithelial cell differentiation, defects in left-right patterning	N/A	Tran et al., 2007, 2010; Kraus et al., 2012; Lian et al., 2014
<i>BMP4</i>	Renal hypodysplasia ⁴	Microphthalmia, syndromic 6, orofacial cleft 11	Important for ventral mesoderm formation. BMP signaling mediates pronephric-tubule and duct formation	Dominant	Bracken et al., 2008; Weber et al., 2008; Neugebauer et al., 2015
<i>BMP7</i>	UVJ obstruction ⁷ , renal hypodysplasia ⁴	CAKUT	Expressed in pronephros	Dominant	Wang et al., 1997; Hwang et al., 2014
<i>CLCNKB</i>	Hypochloremia, proteinuria	Bartter syndrome type III	Important for chloride conductance	Recessive	Simon et al., 1997; Vize, 2003; Seys et al., 2017
<i>EYA1</i>	Renal hypodysplasia ⁴	Branchio-oto-renal syndrome	Important for normal ear development, not analyzed in pronephros	Dominant	Abdelhak et al., 1997; Li et al., 2010
<i>FOXC1</i>	Renal hypoplasia ⁴	CAKUT, Axenfield-Rieger syndrome	Expression in pronephros	Dominant	Nakano et al., 2003; Maguire et al., 2012; Kaminski et al., 2016
<i>FOXC2</i>	Hydronephrosis, horseshoe kidney ⁶ , proteinuria, renal cysts ¹	Lymphedema-distichiasis syndrome with kidney disease and diabetes mellitus	Required for podocyte gene expression	Dominant	Yildirim-Toruner et al., 2004; White et al., 2010
<i>GATA3</i>	Renal dysplasia ¹	HDR syndrome	Expressed in pronephric duct	Dominant	Deconinck et al., 2000; Van Esch et al., 2000
<i>GDNF</i>	Kidney agenesis ³ , renal dysgenesis, nephrotic syndrome	CAKUT, Hirschsprung disease	Expressed in pronephros	N/A	Kyuno and Jones, 2007; Prato et al., 2009
<i>HNF1B</i>	Renal hypodysplasia ⁴ , sporadic renal dysplasia ¹ , renal cysts ¹ , renal agenesis ³ , horseshoe kidney ⁶	MODY5, renal cysts and diabetes syndrome	Distinct renal phenotypes based on specific human mutations	Dominant	Bohn et al., 2003; Gong et al., 2009; Roose et al., 2009; Nagamani et al., 2010; Verdeguer et al., 2010; Drews et al., 2011; Sauert et al., 2012
<i>INVS (NPHP2)</i>	Renal cysts ⁵ , tubular lesions	Nephronophthisis, situs inversus	Role in early tubule morphogenesis	Het and homo	Halbritter et al., 2013; Hoff et al., 2013
<i>LHX1 (XLIM1)</i>	MCDK ¹	CAKUT	Important for proper renal development	N/A	Shawlot and Behringer, 1995; Vize et al., 2003; Nagamani et al., 2010; Drews et al., 2011; DeLay et al., 2018
<i>LMX1B</i>	Renal cysts ¹ , microscopic haematuria, proteinuria	Nail-patella syndrome, nephrotic syndrome, nail-patella-like renal disease	Critical for glomus development, found its downstream position in the gene regulatory network of podocyte specification	Dominant	Miner et al., 2002; Haldin et al., 2008; White et al., 2010; Edwards et al., 2015
<i>NEK8 (NPHP9)</i>	Renal cysts ⁵	Nephronophthisis, Meckel-Gruber syndrome	Role in early tubule morphogenesis	Recessive	Zalli et al., 2012; Hoff et al., 2013
<i>NPHP3</i>	Renal cysts ⁵ , tubular lesions	Nephronophthisis, situs inversus, Meckel-Gruber syndrome, renal-hepatic-pancreatic dysplasia	Role in early tubule morphogenesis	Het and homo	Olbrich et al., 2003; Hoff et al., 2013
<i>NRIP1</i>	Renal hypo/dysplasia ⁴ , VUR ⁷	CAKUT	Important for proper renal development	Dominant	Vivante et al., 2017
<i>PAX2</i>	Renal hypoplasia ⁴ , VUR ⁷ , renal-coloboma syndrome	Pallilorenal syndrome	Essential for tubule differentiation	Dominant	Bouchard et al., 2002; Buisson et al., 2015
<i>PKD1</i>	Renal cysts ¹	ADPKD type 1	Expressed in pronephros	Dominant	Lu et al., 2001; Burtey et al., 2005; Lian et al., 2014
<i>PKD2</i>	Renal cysts ¹	ADPKD type 2	Post-transcriptionally positively regulated by BicC via repression of miR-17	Dominant	Deltas, 2001; Kim et al., 2009; Tran et al., 2010
<i>RET</i>	Renal agenesis ³	CAKUT, Hirschsprung disease	Expressed in pronephric duct	Dominant	Carroll et al., 1998; Skinner et al., 2008

Continued

Table 1. Continued

Gene symbol	Renal phenotype	Human disease	Findings in <i>Xenopus</i>	Mode of inheritance	References
<i>SALL1</i>	Renal hypodysplasia ⁴ , MCDK ¹ , PUV ⁷ , UPJ obstruction ⁷ , VUR ⁷	CAKUT, Townes-Brocks syndrome	Expressed in pronephros	Dominant	Nishinakamura, 2003; Hwang et al., 2014; Kaminski et al., 2016
<i>SIX1</i>	Renal hypodysplasia ⁴	Branchio-oto-renal syndrome	Potential <i>six1</i> gene targets expressed in kidney	Dominant	Ruf et al., 2004; Yan et al., 2015
<i>SIX2</i>	Renal hypodysplasia ⁴	CAKUT	Conserved enhancers regulate pronephros-specific expression	Dominant	Weber et al., 2008; Suzuki et al., 2015
<i>SOX17</i>	VUR ⁷ , UPJ obstruction ⁷	CAKUT	Important for endoderm formation	Dominant	Sinner et al., 2006; Gimelli et al., 2010
<i>TSC1</i>	Kidney tumors ²	Tuberous sclerosis	Downregulated by miRNAs to control proximal tubule size	Dominant	Romaker et al., 2014
<i>WDPCP (FRITZ)</i>	Renal malformation, genital abnormalities	Bardet-Biedel syndrome	Controls septin localization at cilia, linked mutations in human WDPCP to Bardet-Biedel and Meckel-Gruber syndromes	Recessive	Kim et al., 2010a,b; Suspitsin and Imyanitov, 2016
<i>WNT4</i>	Renal hypodysplasia ⁴	Mullerian aplasia and hyperabdrigenism, SERKAL syndrome	Patterns the proximal pronephric primordia, regulated by Notch, Wt1 and Sox11	Dominant	Saulnier et al., 2002; Iglesias et al., 2007; Naylor and Jones, 2009; Murugan et al., 2012; Vivante et al., 2013
<i>WT1</i>	Kidney tumors ²	Wilms-tumor type 1, Denys-Drash syndrome, Frasier syndrome, nephrotic syndrome type 4	Key regulator of glomus development, regulates <i>wnt4</i> expression in the pronephros	Dominant	Kim et al., 2010a,b; White et al., 2010; Murugan et al., 2012

Superscript numbers 1-7 correspond to the numbers found in Fig. 1 (1, green; 2, magenta; 3, purple; 4, teal; 5, orange; 6, blue; 7, red). Bold indicates studies were performed in *Xenopus* to validate that specific mutations cause human kidney phenotypes. CAKUT, congenital anomalies of the kidney and urinary tract; Het, heterozygous; Homo, homozygous; MCDK, multicystic dysplastic kidney; PUV, posterior urethral valves; UPJ, ureteropelvic junction obstruction; UVJ, ureterovesical junction obstruction; VUR, vesicoureteral reflux.

Each segment of the nephron has a specific function, which is reflected by distinct cell morphologies (Møbjerg et al., 2000) and gene expression signatures (Zhou and Vize, 2004; Raciti et al., 2008). Strikingly similar to mammals, the proximal tubules in *Xenopus* are responsible for reabsorbing ions, water, glucose and amino acids (Eid et al., 2002; Zhou and Vize, 2004, 2005; Christensen et al., 2008; Raciti et al., 2008). The most distal part of the *Xenopus* proximal tubule is analogous to the mammalian proximal straight tubule (Raciti et al., 2008). Because *Xenopus* is an aquatic freshwater frog, it is unlikely that the amphibian pronephros requires a true loop of Henle for concentrating urine. Within the literature, the *Xenopus* intermediate tubules have been defined using markers that are co-expressed in the proximal and distal tubules of the mouse kidney (Reggiani et al., 2007; Raciti et al., 2008). Thus, assessment of a marker that is exclusive to this region is necessary to confirm or refute the existence of this structure. The *Xenopus* distal tubule is analogous to the thick ascending limb of the loop of Henle and the distal convoluted tubule in mammals (Raciti et al., 2008); it functions to transport ammonium and reabsorb magnesium ions, and is important for urine acidification (Vize, 2003; Zhou and Vize, 2004, 2005; Raciti et al., 2008). The pronephric connecting tubule of *Xenopus* connects the distal tubules to the cloaca to excrete urine and is only analogous to mammals in the region neighboring the distal tubule (Raciti et al., 2008). The pronephros does not have a collecting duct to further concentrate urine or connect multiple nephrons to a secondary structure, as is necessary for the mammalian metanephros.

Recent research has shown that human and mouse nephron protein-expression signatures in the renal vesicles (Box 1) and S-shaped bodies are very similar (Lindström et al., 2018). Not surprisingly, *Xenopus* shares similar gene expression signatures to humans during early pronephric development in the kidney (Fig. 3). Although Fig. 3 only illustrates four gene expression signatures, it is

important to note that, of the 29 expression signatures identified in the human embryonic kidney (Lindström et al., 2018), 18 can also be found in the *Xenopus* kidney, including *mafb*, *slc3a1*, *pou3f3* and *foxc2* (xenbase.org). Additionally, a comparison of gross expression signatures in the kidney demonstrated similarities between mouse and *Xenopus* in both overlapping transcription factors such as *Pax8* and additional ones such as *Foxc1* and *Sall1* (Kaminski et al., 2016). Furthermore, some of the proteins analyzed in humans have not been studied in *Xenopus*, potentially leading to an underestimate of the similarities. This overall conservation on the molecular, genetic, structural and functional levels supports the use of the *Xenopus* pronephros as an appropriate model for the study of CAKUT.

Delineating molecular pathways involved in kidney development using *Xenopus*

Xenopus has been historically used to elucidate molecular mechanisms and signaling cascades involved in early developmental processes (Beck and Slack, 2001). More recently, *Xenopus* has played an important role in identifying the function of genes that are involved in CAKUT. The Wnt pathway has been heavily studied in *Xenopus* due to this pathway's significant contribution to various developmental processes. The canonical Wnt/β-catenin pathway plays a role in kidney development in both mice (Carroll et al., 2005; Iglesias et al., 2007) and *Xenopus* (Saulnier et al., 2002; Lyons et al., 2009). WNT4 is a well-known ligand that activates canonical Wnt signaling and is important for induction and mesenchyme-to-epithelial transition of the metanephric mesenchyme (Box 1) (Murugan et al., 2012). In humans, *WNT4* loss-of-function mutations have been associated with renal hypodysplasia, a condition characterized by dysplastic kidneys with a reduced number of nephrons (Vivante et al., 2013). Studies in *Xenopus* demonstrate that Wnt4 protein controls the medio-lateral patterning of the pronephros (Naylor and Jones, 2009).




Fig. 3. Expression patterns are conserved between the human S-shaped body and the early *Xenopus* pronephros during development. Four different expression patterns (shown in different colors) of fundamental kidney proteins/mRNA were chosen to demonstrate the similarities between the developing human and *Xenopus* nephron. Schematics indicate where immunostaining of S-shaped body nephrons of week-16 to -17 human fetal kidneys is present (top) (Lindström et al., 2018), as well as *in situ* expression patterns of stage-33 *Xenopus* nephrons (bottom) (Xenbase.org). Schematics are positioned so that the proximal and distal regions of the human and *Xenopus* nephron expression patterns can be easily compared. The *Xenopus* kidney stage was chosen to match the approximate developmental time point of human S-shaped body nephrons, as both are representative of recently epithelialized nephrons. Note that *pax2* is slightly more enriched at the nephrostomes in *Xenopus* (dark green).

Additionally, knockdown of Wnt4 protein in *Xenopus* results in complete loss of kidney tubules (Saulnier et al., 2002).

In addition to Wnt ligands, other proteins have been found to regulate Wnt signaling. In humans, Wilms tumor 1 (*WT1*) is a known regulator of canonical Wnt signaling. Mutations that affect *WT1* result in nephroblastoma, more commonly known as Wilms tumor (Kim et al., 2010a,b). Knockdown of Wt1 protein expression in *Xenopus* embryos was shown to reduce Wnt4 expression in the prospective pronephros (Murugan et al., 2012). Mutations in the RNA-binding protein BICC1, a negative Wnt regulator, result in a cystic kidney phenotype in mice and renal anomalies in patients (Kraus et al., 2012). Prior studies in *Xenopus* indicate that *Bicc1* inhibits the microRNA miR-17, preventing miR-17 from destabilizing the *pkd2* transcript (Tran et al., 2010). This was demonstrated by a rescue of the mutant kidney phenotype when both miR-17 and *Bicc1* were simultaneously knocked down in tadpoles (Tran et al., 2010). Interestingly, loss of the *Pkd1* gene product, polycystin-1, downregulates BICC1 expression in mouse kidneys and mouse cell lines (Lian et al., 2014). The connection between BICC1 downregulation and polycystin-1 loss in mice and the stabilization of *pkd2* mRNA by *Bicc1* in *Xenopus* suggests that disruption of BICC1 may induce cystic phenotypes through polycystin signaling (Box 1). Similarly, targeted deletion of *Hnf1b*, a key transcription factor involved in kidney development, decreases *Bicc1* and *Pkd2* mRNA expression in the mouse kidney (Verdeguer et al., 2010). HNF1B in the mouse (Gresh et al., 2004; Verdeguer et al., 2010) and BICC1 in mice and *Xenopus* (Tran et al., 2010) regulate the *Pkd2* transcript, and mutations in either gene can lead to kidney cysts in mouse models (Gong et al., 2009; Tran et al., 2010). Thus, it is likely that these genes act through similar pathways. *Xenopus* would be an ideal model for studying the components of this pathway because molecular strategies to assess them have already been established (Tran et al., 2007, 2010).

Another common congenital kidney disease whose pathway has been elucidated using *Xenopus* is nephronophthisis (NPHP; Box 1). NPHP is a ciliopathy (Box 1), resulting in either abnormal formation or function of cilia (Box 1). NPHP is a rare birth defect, but it is the most common cause of kidney failure, or end-stage renal disease (ESRD; Box 1), in the first three decades of life (Wolf and Hildebrandt, 2011). Studies in *Xenopus* linked several genes with NPHP or clarified components of the mechanism that causes the disease. First, targeted kidney knockdown of the Wnt signaling inhibitor *Invs* (inversin; also known as NPHP2) demonstrated that

Invs is important for morphogenetic cell movements during tubule elongation (Lienkamp et al., 2010). This knockdown led to impaired ventral proximal pronephros extension and distal tubule differentiation (Lienkamp et al., 2010). Later, it was discovered that several genes that cause NPHP form a distinct complex that contributes to kidney development. In *Xenopus*, *Anks6* was identified as an NPHP family member that assembles a protein complex of the NPHP-associated proteins Nek8, *Invs* and *Nphp3* to regulate kidney development (Hoff et al., 2013). This study showed that *Anks6* localizes to the base of the cilium and that knockdown of its protein expression results in kidney anomalies (Hoff et al., 2013). The study also linked *ANKS6* mutations in patients with an NPHP-like clinical syndrome (Hoff et al., 2013). Additionally, knockdown of either *Anks6* or *Nphp3* protein in *Xenopus* embryos results in edema, which suggests that they may be important for kidney function (Hoff et al., 2013). Likewise, edema has been shown to occur due to depletion of *Invs* protein (Lienkamp et al., 2010).

Although *Xenopus* has helped uncover some of the causative genes for NPHP and their role in kidney development and function, only ~40% of patients have a mutation in one of the 20 known NPHP-related genes (Stokman et al., 2016; Srivastava et al., 2018). A recent genome-wide association study (GWAS; Box 1) has uncovered epigenetic signatures of CKD (Box 1), including *NPHP4* (Wing et al., 2014). Some NPHP patients may suffer from epigenetic alterations of NPHP genes rather than a mutation in the gene itself. Future research using *Xenopus* may lead to the identification of new NPHP-causing genes and may uncover further epigenetic regulation of known genes as they have been used to study epigenetics.

A notable CAKUT-causing gene encodes the transcription factor *LMX1B*, mutations in which can lead to nail-patella syndrome (NPS; Box 1), a rare cause of autosomal-dominant ESRD (Edwards et al., 2015). Additionally, mutations in *LMX1B* can result in glomerulopathies (Box 1). Mouse studies suggest that *Lmx1b* is necessary for the maintenance of podocytes, which are highly specialized cells that wrap around capillaries of the glomerulus and restrict the passing of macromolecules into the kidney (Miner et al., 2002). In *Xenopus*, *lmx1b* is expressed in the glomus, which is analogous to the mammalian glomerulus, and is also known to have a role in podocyte specification (Haldin et al., 2008; White et al., 2010). Morpholino knockdown of *Lmx1b* protein in *Xenopus* showed that the glomus was reduced in size and the development of the proximal tubules was limited (Haldin et al., 2008). Taken

together, these studies demonstrate the usefulness of *Xenopus* in uncovering the role of proteins important for kidney development and function. Additionally, they show that *Xenopus* is capable of recapitulating specific kidney disease phenotypes found in humans.

Using *Xenopus* to examine human genetic variants

In human studies, missense mutations are often analyzed via software programs such as PredictSNP (Bendl et al., 2014) and Meta-SNP (Capriotti et al., 2013). These programs predict the functional relevance of an amino acid change and suggest whether they are likely to be pathogenic. One drawback of this method is the surprising data from the 1000 Genomes Project indicating that people with predicted pathogenic variants do not suffer from the expected disease (Xue et al., 2012). *Xenopus* can be used to verify the pathogenicity and expressivity (Box 1) of potential disease-causing genetic variants *in vivo*.

Work in mouse models found that *CLCNKB* is important for concentrating urine (Matsumura et al., 1999). In humans, *CLCNKB* mutations cause a salt-losing tubulopathy (Box 1) known as Bartter syndrome type III (Simon et al., 1997). The first genotype-phenotype correlation for Bartter syndrome type III was validated by functional analysis of eight missense and two nonsense mutations in *Xenopus*. By performing voltage-clamp experiments in *Xenopus* oocytes, it was found that nine of the ten mutations significantly decreased normal conductance (Seys et al., 2017), indicating altered salt homeostasis. This work verified that the reported human genetic mutations were, in fact, Bartter syndrome type III-causing mutations (Seys et al., 2017). A similar study demonstrated that dysregulation of NRIP1-dependent retinoic acid signaling in both *Xenopus* and mouse disrupted kidney formation (Vivante et al., 2017). In *Xenopus*, knockdown of *Nrip1* protein causes kidney anomalies that can be rescued with wild-type human *NRIP1* mRNA but not with the truncated *NRIP1* mRNA identified in affected individuals (Vivante et al., 2017). *Xenopus* can also serve to assess the distinct effects of different mutations in a single gene.

HNF1B mutations have been shown to manifest as distinct renal diseases in humans (Bohn et al., 2003). An attempt to model these differences in *Xenopus* was successfully executed using different methods in two separate studies. In one study, researchers generated transgenic *Xenopus* lines expressing two different human *HNF1B* mutants using a heat-shock-inducible (Box 1) Cre-*loxP* system (Sauert et al., 2012). One transgenic line expressed an insertion mutation, while the other line expressed a deletion. The deletion led to reduced pronephric development, while the insertion enlarged the pronephros, with both phenotypes primarily affecting the proximal tubules (Sauert et al., 2012). In another study, Bohn and colleagues compared nine different human *HNF1B* mutations, including indels, missense and nonsense mutations, by injecting mutant mRNA. The mutations led to distinct renal disease phenotypes in *Xenopus*, as they do in humans (Bohn et al., 2003). In *Xenopus*, six of the mutants resulted in an enlargement of the pronephric structures, while the other three mutations led to a reduction or loss of the tubules and the anterior part of the duct (Bohn et al., 2003). The range of kidney phenotypes seen in *Xenopus* recapitulate the kidney phenotypes observed in humans. This potentially allows for a correlation between an observed patient phenotype with a specific mutation in the *HNF1B* gene. Studies like these are an efficient way to identify pathogenic genetic variants and can represent the variation of expressivity seen in patients.

Early embryonic lethality

In *Xenopus*, many genes that are involved in kidney development have essential functions in other tissues. To bypass embryonic

lethality due to the loss of these essential proteins, researchers have developed techniques that are not possible in other organisms. For example, tissue-targeted microinjection has been used to study ADPKD, the most common inherited human renal disease. ADPKD is caused by mutations in either *PKD1* or *PKD2*, with *PKD2* mutations causing a less severe form of ADPKD with a later onset than *PKD1* mutations (Deltas, 2001). Homozygous mutations or deletions of either *Pkd1* or *Pkd2* in mouse models are embryonic lethal (Lu et al., 2001; Kim et al., 2009). This limitation can be resolved by tissue-targeted microinjections in *Xenopus* as described earlier, as both *pkd1* (Burtey et al., 2005) and *pkd2* (Tran et al., 2010) are expressed in the *Xenopus* embryonic kidney. Using this technique, Tran and colleagues elucidated some aspects of *pkd2* post-transcriptional regulation (Tran et al., 2010). Although a variety of mouse models have since been developed to bypass the embryonic lethality of a complete knockout mouse, there is still a need for ADPKD models that more closely resemble the human disease. *Xenopus* has a lifespan that permits an extended survival that would add additional information of how ADPKD progresses over time. Additionally, elucidating the mechanism of the disease using *Xenopus* may be more cost and time effective.

Inducible systems have also been successfully used in *Xenopus* embryos, which allows for temporal or spatial control of gene expression. Four inducible systems were developed in *Xenopus*: the GAL4-UAS (Box 1) (Denayer et al., 2006), Tet-On (Box 1) (Rankin et al., 2011), heat-shock inducible (Roose et al., 2009) and the Dex-inducible (Box 1) strategy (Zhuo et al., 2013). In addition, knocking down a specific protein can be titrated, either by using morpholinos at varying doses to reduce lethality (Eisen and Smith, 2008) or by microinjecting less mRNA of an inhibitor of the protein of interest (Baker et al., 1999). Alternatively, the CRISPR/Cas9 system can be utilized in *Xenopus* to bypass early developmental lethality caused by knocking out essential genes such as *lhx1* (DeLay et al., 2018). Human 17q12 deletions, which span two genes important for kidney development, *HNF1B* and *LHX1*, have been implicated in CAKUT (Nagamani et al., 2010). Although embryonically lethal in mice, *Xenopus* embryos survive *lhx1* knockout and demonstrate abnormal kidney development (DeLay et al., 2018). CRISPR/Cas9 editing results in mosaic knockout in *Xenopus*, which may explain why the knockout embryos were able to bypass embryonic lethality (DeLay et al., 2018), making this mosaic expression a silver lining. Furthermore, in *Xenopus tropicalis*, transplanting the primordial germ cells of CRISPR-mutagenized F₀ embryos into a wild-type host, termed leapfrogging (Box 1), results in F₁ embryos bypassing the embryonic lethality of the loss of the homeobox protein gooseoid (Blitz et al., 2016). Taken together, there are a range of techniques available to bypass embryonic lethality in *Xenopus*.

Versatility in understanding cilia formation

Cilia are hair-like cellular protrusions that are found in most tissues in the body. Defects in genes important for ciliogenesis often lead to a vast array of phenotypes, including retinal degeneration, renal disease and congenital fibrocystic diseases of the liver (Waters and Beales, 2011). In addition to studying the ciliated A6 cell line derived from *Xenopus* and the primary cilia present *in vivo* in the pronephros, *Xenopus* embryos have a ciliated epidermis that has been useful for studying cilia function. *WDPCP* (Fritz) is one of 20 known genes that cause Bardet-Biedl syndrome (BBS; Box 1), all of which are involved in primary cilia function (Suspitsin and Imyanitov, 2016). Primary features of BBS include rod-cone dystrophy, genital abnormalities, learning difficulties, obesity and renal defects (Suspitsin and Imyanitov, 2016). The *Xenopus* epidermis has been

used to study BBS because the cilia are easy to visualize. The authors found that *Wdpcp* localizes cytoskeletal proteins that control cell movement, called septins (Box 1), to the ciliary base (Kim et al., 2010a,b). Additionally, mutations in human *WDPCP* are linked to Meckel-Gruber (Box 1) syndrome (Kim et al., 2010a,b). This study demonstrates the versatility that the *Xenopus* model provides in understanding kidney diseases that extend beyond the pronephros.

Future approaches

Until now, many studies utilized morpholinos to knock down protein expression. Additionally, the allotetraploid genome and long generation time (1 year) of *X. laevis* have previously made it less attractive for genetic experiments than the diploid *X. tropicalis* (Beck and Slack, 2001). However, the field is moving towards using CRISPR/Cas9 technology to generate gene knockouts. CRISPR/Cas9 gene editing in *Xenopus* is a rapid and inexpensive means of studying genes involved in developmental processes (Bhattacharya et al., 2015), introducing the possibility of performing large-scale genetic screens. Recently, the CRISPR/Cas9 system has been used in *Xenopus* to knock out an important kidney gene, *lhx1* (DeLay et al., 2018). Knock-in strategies using CRISPR/Cas9 have also been successfully utilized in *X. tropicalis* (Shi et al., 2015). Targeted CRISPR/Cas9 knock-in strategies facilitate the fusion of fluorescent tags to endogenous genes (Shi et al., 2015), allowing for more precise visualization of proteins *in vivo*. Furthermore, leapfrogging, which effectively circumvents a generation of breeding, allows researchers to generate transgenic lines quicker than with traditional breeding strategies (Blitz et al., 2016). These technologies are being favored over new criticisms concerning morpholino-driven phenotypes. Morpholinos have been recently shown to cause immune-response-related gene transcription and mistargeted splicing events. Although these effects cannot be completely ameliorated, researchers can take steps to reduce them, such as lowering the GC content and titrating the morpholino (Gentsch et al., 2018). Previous studies using morpholinos should not be discarded though, as our group's recent work suggests that they match CRISPR/Cas9 phenotypes (DeLay et al., 2019), but optimization to reduce unintended effects as well as proper controls should be utilized.

Distinctly, *Xenopus* has been used to study acute kidney injury and regeneration. Upon unilateral nephrectomy of the proximal tubules at stage 37/38 (~53 hpf), *Xenopus* tadpoles are able to regenerate functional proximal tubules with no drug intervention (Caine and McLaughlin, 2013; Suzuki et al., 2019). Additionally, *Xenopus* can even be used to generate kidney organoids that can replace the *Xenopus* kidney *in vivo* (Chan et al., 1999), which broadens the usefulness of this model organism for studying kidney disease. Because *Xenopus* oocytes are equipped with all the necessary machinery for development upon fertilization, they have long been used to study channel proteins. Given the large number of ion channels present in the kidney, this system can be readily exploited to study patient mutations in various channels and their effects on conductance. Additionally, two new transgenic frog lines have been developed in which GFP is expressed in the developing pronephros and mesonephros (Corkins et al., 2018), as well as in the pronephros alone (Ochi et al., 2012). This permits live imaging of the developing kidney, which can also be used to assess kidney disruption upon specific gene knockdown/out without the need for immunostaining or *in situ* hybridization. This new *Xenopus* line has the potential to streamline large-scale screens of kidney-disease-causing genes and of environmental factors that affect renal development. Also, bypassing the need to process embryos will expedite drug discovery screening in *Xenopus* (Schmitt et al., 2014), which may ultimately lead to

clinical applications. The development of these new tools will undoubtedly advance our ability to study the genetic mechanisms underlying kidney development and disease.

Limitations of *Xenopus* as a kidney model

Although *Xenopus* has many advantages that make it an attractive model for kidney development and disease, like all other models it possesses its own set of limitations. The most notable limitation may be that *Xenopus* is used to study the pronephros and has mesonephric nephrons that serve as its adult kidney instead of the metanephros of mammals. Although these differing nephrons maintain the same structure and function, this should be taken into consideration. Studying the pronephros may be different, but it offers a simplified method to study non-integrated individual nephrons. Many cases of CAKUT are associated with structural developmental defects such as vesicoureteral reflux, which cannot be visualized in *Xenopus* as they do not have a ureteric bud that forms a true collecting-duct system. Therefore, for malformations such as this, researchers should study *Xenopus* as a way to implicate new proteins in CAKUT and use in concurrence with another model system, such as mouse or kidney organoids, to study the anomalies more directly. Additionally, the *Xenopus* glomus does not have a Bowman's space, but instead blood filtrate goes directly into the coelomic cavity. Although this should be considered when conducting studies concerning the glomerulus, it may have a separate advantage. In mice, the glomerulus forms at the most proximal end of the S-shaped body, linking it to the development of the rest of the nephron. *Xenopus* however, can develop a glomus independently from the tubules, which may make understanding how the glomus forms more accessible (Urban et al., 2006). Lastly, *Xenopus* is an aquatic organism that does not require the resorption of water that is necessary in terrestrial animals.

Conclusion

The studies reviewed here demonstrate that *Xenopus* is well suited to model congenital kidney diseases. *Xenopus* shares many genes and proteins with the mammalian kidney that have similar expression patterns and functions. As described here, many of the renal phenotypes observed in humans can be recapitulated in frogs. Next-generation DNA sequencing strategies, such as whole-exome sequencing, has helped identify candidate CAKUT-causing genes. An ideal organism to functionally assay these genes *in vivo* would develop quickly and produce large numbers of embryos, qualities that *Xenopus* possesses.

CAKUT accounts for 40-50% of pediatric chronic kidney disease worldwide (Vivante et al., 2014). Although important advances have been made in basic nephrology research, there are still missing pieces in the molecular basis of kidney development. This failure to fully grasp how the kidney develops translates into low cure rates for many of the genetic disorders that cause kidney disease. Being able to uncover novel genes and pathways that are involved in nephrogenesis will give us a better understanding of kidney development as a whole. This may lead to better treatment for patients with CAKUT, and its causative mutations may ultimately be eliminated with the application of gene therapy in human embryos. *Xenopus* is a very promising organism that can get us closer to accomplishing these goals.

Acknowledgements

We thank Richard Behringer, Mark Corkins, Bridget DeLay and Vanja Krmeta-Stankic for critical reading of the manuscript and thoughtful suggestions.

Competing interests

The authors declare no competing or financial interests.

Author contributions

A.T.M.B. wrote the Review and created the figures. R.K.M. provided conceptual input into the writing and figures.

Funding

This work was funded by National Institutes of Health NIDDK grants (K01DK092320 and R03 DK118771) and startup funding from the Department of Pediatrics at the University of Texas Health Science Center at Houston's McGovern Medical School.

References

- Abdelhak, S., Kalatzis, V., Heilig, R., Compain, S., Samson, D., Vincent, C., Weil, D., Cruaud, C., Sahly, I., Leibovici, M. et al. (1997). A human homologue of the *Drosophila* eyes absent gene underlies branchio-oto-renal (BOR) syndrome and identifies a novel gene family. *Nat. Genet.* **15**, 157-164.
- Al-Maawali, A., Walfisch, A. and Koren, G. (2012). Taking angiotensin-converting enzyme inhibitors during pregnancy: is it safe? *Can. Fam. Physician* **58**, 49-51.
- Aslan, Y., Tadjuidje, E., Zorn, A. M. and Cha, S.-W. (2017). High efficiency non-mosaic CRISPR mediated knock-in and mutations in F0 *Xenopus*. *Development* **144**, 2852-2858.
- Baker, J. C., Beddington, R. S. P. and Harland, R. M. (1999). Wnt signaling in *Xenopus* embryos inhibits *Bmp4* expression and activates neural development. *Genes Dev.* **13**, 3149-3159.
- Beck, C. W. and Slack, J. M. (2001). An amphibian with ambition: a new role for *Xenopus* in the 21st century. *Genome Biol.* **2**, REVIEWS1029.
- Bend, J., Stourac, J., Salanda, O., Pavelka, A., Wieben, E. D., Zendulka, J., Brezovsky, J. and Damborsky, J. (2014). PredictSNP: robust and accurate consensus classifier for prediction of disease-related mutations. *PLoS Comput. Biol.* **10**, e1003440.
- Bertram, J. F., Douglas-Denton, R. N., Diouf, B., Hughson, M. D. and Hoy, W. E. (2011). Human nephron number: Implications for health and disease. *Pediatr. Nephrol.* **26**, 1529-1533.
- Bhattacharya, D., Marfo, C. A., Li, D., Lane, M. and Khokha, M. K. (2015). CRISPR/Cas9: An inexpensive, efficient loss of function tool to screen human disease genes in *Xenopus*. *Dev. Biol.* **408**, 196-204.
- Black, M. J., Sutherland, M. R. and Gubhaju, L. (2012). Effects of preterm birth on the kidney. In *Basic Nephrology and Acute Kidney Injury* (Ed. Prof. Manisha Sahay), pp. 61-88. Intech.
- Blitz, I. L., Fish, M. B. and Cho, K. W. Y. (2016). Leapfrogging: primordial germ cell transplantation permits recovery of CRISPR/Cas9-induced mutations in essential genes. *Development* **143**, 2868-2875.
- Bohn, S., Thomas, H., Turan, G., Ellard, S., Bingham, C., Hattersley, A. T. and Ryffel, G. U. (2003). Distinct molecular and morphogenetic properties of mutations in the human *HNF1beta* gene that lead to defective kidney development. *J. Am. Soc. Nephrol.* **14**, 2033-2041.
- Boualila, S. K., Gaitan, Y., Tremblay, M., Sharma, R., Cardin, J., Kania, A. and Bouchard, M. (2013). A core transcriptional network composed of *Pax2/8*, *Gata3* and *Lim1* regulates key players of pro/mesonephros morphogenesis. *Dev. Biol.* **382**, 555-566.
- Bouchard, M., Souabni, A., Mandler, M., Neubüser, A. and Busslinger, M. (2002). Nephric lineage specification by *Pax2* and *Pax8*. *Genes Dev.* **16**, 2958-2970.
- Bracken, C. M., Mizeracka, K. and McLaughlin, K. A. (2008). Patterning the embryonic kidney: BMP signaling mediates the differentiation of the pronephric tubules and duct in *Xenopus laevis*. *Dev. Dyn.* **237**, 132-144.
- Buisson, I., Le Bouffant, R., Futel, M., Riou, J. F. and Umbhauer, M. (2015). *Pax8* and *Pax2* are specifically required at different steps of *Xenopus* pronephros development. *Dev. Biol.* **397**, 175-190.
- Burtey, S., Leclerc, C., Nabais, E., Munch, P., Gohory, C., Moreau, M. and Fontés, M. (2005). Cloning and expression of the amphibian homologue of the human *PKD1* gene. *Gene* **357**, 29-36.
- Caine, S. T. and McLaughlin, K. A. (2013). Regeneration of functional pronephric proximal tubules after partial nephrectomy in *Xenopus laevis*. *Dev. Dyn.* **242**, 219-229.
- Capriotti, E., Altman, R. B. and Bromberg, Y. (2013). Collective judgment predicts disease-associated single nucleotide variants. *BMC Genomics* **14** Suppl. 3, S2.
- Carroll, T., Wallingford, J. B., Seufert, D. and Vize, P. D. (1998). Molecular regulation of pronephric development. *Curr. Top. Dev. Biol.* **44**, 67-100.
- Carroll, T. J., Park, J. S., Hayashi, S., Majumdar, A. and McMahon, A. P. (2005). *Wnt9b* plays a central role in the regulation of mesenchymal to epithelial transitions underlying organogenesis of the mammalian urogenital system. *Dev. Cell* **9**, 283-292.
- Chan, T.-C., Ariizumi, T. and Asashima, M. (1999). A model system for organ engineering: Transplantation of in vitro induced embryonic kidney. *Naturwissenschaften* **86**, 224-227.
- Christensen, E. I., Raciti, D., Reggiani, L., Verroust, P. J. and Brändli, A. W. (2008). Gene expression analysis defines the proximal tubule as the compartment for endocytic receptor-mediated uptake in the *Xenopus* pronephric kidney. *Pflugers Arch.* **456**, 1163-1176.
- Corkins, M. E., Hanania, H. L., Krneta-Stankic, V., DeLay, B. D., Pearl, E. J., Lee, M., Ji, H., Davidson, A. J., Horb, M. E. and Miller, R. K. (2018). Transgenic *Xenopus laevis* line for in vivo labeling of nephrons within the kidney. *Genes* **9**, 197.
- Costantini, F. and Kopan, R. (2010). Patterning a complex organ: Branching morphogenesis and nephron segmentation in kidney development. *Dev. Cell*, 698-712.
- Dale, L. and Slack, J. M. W. (1987). Fate map for the 32-cell stage of *Xenopus laevis*. *Development* **99**, 527-551.
- Davidson, A. (2008). Mouse kidney development. *StemBook*.
- Deconinck, A. E., Mead, P. E., Tevosian, S. G., Crispino, J. D., Katz, S. G., Zon, L. I. and Orkin, S. H. (2000). FOG acts as a repressor of red blood cell development in *Xenopus*. *Development* **127**, 2031-2040.
- DeLay, B. D., Krneta-Stankic, V. and Miller, R. K. (2016). Technique to target microinjection to the developing *xenopus* kidney. *J. Vis. Exp.* **111**, e53799.
- DeLay, B. D., Corkins, M. E., Hanania, H. L., Salanga, M., Deng, J. M., Sudou, N., Taira, M., Horb, M. E. and Miller, R. K. (2018). Tissue-specific Gene Inactivation in *Xenopus laevis*: Knockout of *lhx1* in the Kidney with CRISPR/Cas9. *Genetics* **208**, 673-686.
- DeLay, B. D., Baldwin, T. A. and Miller, R. K. (2019). Dynamins binding protein is required for *Xenopus laevis* kidney development. *Front. Physiol.* **10**, R8632.
- Deltas, C. C. (2001). Mutations of the human polycystic kidney disease 2 (PKD2) gene. *Hum. Mutat.* **18**, 13-24.
- Denayer, T., Van Roy, F. and Vleminckx, K. (2006). In vivo tracing of canonical Wnt signaling in *Xenopus* tadpoles by means of an inducible transgenic reporter tool. *FEBS Lett.* **580**, 393-398.
- Dressler, G. R. (2006). The cellular basis of kidney development. *Annu. Rev. Cell Dev. Biol.* **22**, 509-529.
- Drews, C., Senkel, S. and Ryffel, G. U. (2011). The nephrogenic potential of the transcription factors *osr1*, *osr2*, *hnf1b*, *lhx1* and *pax8* assessed in *Xenopus* animal caps. *BMC Dev. Biol.* **11**, 5.
- Edwards, N., Rice, S. J., Raman, S., Hynes, A. M., Srivastava, S., Moore, I., Al-Hamed, M., Xu, Y., Santibanez-Koref, M., Thwaites, D. T. et al. (2015). A novel *LMX1B* mutation in a family with end-stage renal disease of "unknown cause". *Clin. Kidney J.* **8**, 113-119.
- Eid, S. R., Terretaz, A., Nagata, K. and Brändli, A. W. (2002). Embryonic expression of *Xenopus* SGLT-1L, a novel member of the solute carrier family 5 (SLC5), is confined to tubules of the pronephric kidney. *Int. J. Dev. Biol.* **46**, 177-184.
- Eisen, J. S. and Smith, J. C. (2008). Controlling morpholino experiments: don't stop making antisense. *Development* **135**, 1735-1743.
- Fox, H. (1963). The Amphibian Pronephros. *Q. Rev. Biol.* **38**, 1-25.
- Gentsch, G. E., Spruce, T., Monteiro, R. S., Owens, N. D. L., Martin, S. R. and Smith, J. C. (2018). Innate immune response and off-target mis-splicing are common morpholino-induced side effects in *Xenopus*. *Dev. Cell* **44**, 597-610.e10.
- Gimelli, S., Caridi, G., Beri, S., McCracken, K., Boccia, R., Zordan, P., Dagnino, M., Fiorio, P., Murer, L., Benetti, E. et al. (2010). Mutations in *SOX17* are associated with congenital anomalies of the kidney and the urinary tract. *Hum. Mutat.* **31**, 1352-1359.
- Gong, Y., Ma, Z., Patel, V., Fischer, E., Hiesberger, T., Pontoglio, M. and Igarashi, P. (2009). *HNF-1* regulates transcription of the PKD modifier gene *Kif12*. *J. Am. Soc. Nephrol.* **20**, 41-47.
- Gresh, L., Fischer, E., Reimann, A., Tanguy, M., Garbay, S., Shao, X., Hiesberger, T., Fiette, L., Igarashi, P., Yaniv, M. et al. (2004). A transcriptional network in polycystic kidney disease. *EMBO J.* **23**, 1657-1668.
- Gribouval, O., Morinière, V., Pawtowski, A., Arrondel, C., Sallinen, S. L., Saloranta, C., Clericuzio, C., Viot, G., Tantau, J., Blesson, S. et al. (2012). Spectrum of mutations in the renin-angiotensin system genes in autosomal recessive renal tubular dysgenesis. *Hum. Mutat.* **33**, 316-326.
- Halbritter, J., Porath, J. D., Diaz, K. A., Braun, D. A., Kohl, S., Chaki, M., Allen, S. J., Soliman, N. A., Hildebrandt, F. and Otto, E. A. (2013). Identification of 99 novel mutations in a worldwide cohort of 1,056 patients with a nephronophthisis-related ciliopathy. *Hum. Genet.* **132**, 865-884.
- Haldin, C. E., Massé, K. L., Bhamra, S., Simrick, S., Kyuno, J.-I. and Jones, E. A. (2008). The *lhx1b* gene is pivotal in glomus development in *Xenopus laevis*. *Dev. Biol.* **322**, 74-85.
- Halt, K. and Vainio, S. (2014). Coordination of kidney organogenesis by Wnt signaling. *Pediatr. Nephrol.* **29**, 737-744.
- Hellsten, U., Harland, R. M., Gilchrist, M. J., Hendrix, D., Jurka, J., Kapitonov, V., Ovcharenko, I., Putnam, N. H., Shu, S., Taher, L. et al. (2010). The genome of the Western clawed frog *Xenopus tropicalis*. *Science* **328**, 633-636.
- Hoff, S., Halbritter, J., Epting, D., Frank, V., Nguyen, T.-M. T., van Reeuwijk, J., Boehlke, C., Schell, C., Yasunaga, T., Helmstädter, M. et al. (2013). ANKS6 is a central component of a nephronophthisis module linking NEK8 to INVS and NPHP3. *Nat. Genet.* **45**, 951-956.
- Holbrook, J. D. and Danpure, C. J. (2002). Molecular basis for the dual mitochondrial and cytosolic localization of alanine:glyoxylate aminotransferase in amphibian liver cells. *J. Biol. Chem.* **277**, 2336-2344.

- Hurtado del Pozo, C., Garreta, E., Izpisua Belmonte, J. C. and Montserrat, N. (2018). Modeling epigenetic modifications in renal development and disease with organoids and genome editing. *Dis. Model. Mech.* **11**, dmm035048.
- Hwang, D. Y., Dworschak, G. C., Kohl, S., Saisawat, P., Vivante, A., Hilger, A. C., Reutter, H. M., Soliman, N. A., Bogdanovic, R., Kehinde, E. O. et al. (2014). Mutations in 12 known dominant disease-causing genes clarify many congenital anomalies of the kidney and urinary tract. *Kidney Int.* **85**, 1429-1433.
- Iglesias, D. M., Hueber, P.-A., Chu, L., Campbell, R., Patenaude, A.-M., Dziarmaga, A. J., Quinlan, J., Mohamed, O., Dufort, D. and Goodyer, P. R. (2007). Canonical WNT signaling during kidney development. *Am. J. Physiol. Renal Physiol.* **293**, F494-F500.
- Kaminski, M. M., Tosic, J., Kresbach, C., Engel, H., Klockenbusch, J., Müller, A.-L., Pichler, R., Grahamer, F., Kretz, O., Huber, T. B. et al. (2016). Direct reprogramming of fibroblasts into renal tubular epithelial cells by defined transcription factors. *Nat. Cell Biol.* **18**, 1269-1280.
- Kim, I., Ding, T., Fu, Y., Li, C., Cui, L., Li, A., Lian, P., Liang, D., Wand, D. W., Guo, C. et al. (2009). Conditional mutation of Pkd2 causes cystogenesis and upregulates β -catenin. *J. Am. Soc. Nephrol.* **20**, 2556-2569.
- Kim, M. S., Yoon, S. K., Bollig, F., Kitagaki, J., Hur, W., Whye, N. J., Wu, Y. P., Rivera, M. N., Park, J. Y., Kim, H. S. et al. (2010a). A novel wilms tumor 1 (WT1) target gene negatively regulates the WNT signaling pathway. *J. Biol. Chem.* **285**, 14585-14593.
- Kim, S. K., Shindo, A., Park, T. J., Oh, E. C., Ghosh, S., Gray, R. S., Lewis, R. A., Johnson, C. A., Attie-Bittach, T., Katsanis, N. et al. (2010b). Planar cell polarity acts through septins to control collective cell movement and ciliogenesis. *Science* **329**, 1337-1340.
- Kraus, M. R. C., Clauin, S., Pfister, Y., Di Maio, M., Ulinski, T., Constam, D., Bellanné-Chantelot, C. and Grapin-Botton, A. (2012). Two mutations in human BICC1 resulting in wnt pathway hyperactivity associated with cystic renal dysplasia. *Hum. Mutat.* **33**, 86-90.
- Kyuno, J. and Jones, E. A. (2007). GDNF expression during Xenopus development. *Gene Expr. Patterns* **7**, 313-317.
- Li, Y., Manaligod, J. M. and Weeks, D. L. (2010). EYA1 mutations associated with the branchio-oto-renal syndrome result in defective otic development in *Xenopus laevis*. *Biol. Cell* **102**, 277-292.
- Lian, P., Li, A., Li, Y., Liu, H., Liang, D., Hu, B., Lin, D., Jiang, T., Moeckel, G., Qin, D. et al. (2014). Loss of polycystin-1 inhibits Bicc1 expression during mouse development. *PLoS ONE* **9**, e88816.
- Lienkamp, S., Ganner, A., Boehlke, C., Schmidt, T., Arnold, S. J., Schäfer, T., Romaker, D., Schuler, J., Hoff, S., Powelske, C. et al. (2010). Inversin relays Frizzled-8 signals to promote proximal pronephros development. *Proc. Natl. Acad. Sci. USA* **107**, 20388-20393.
- Lindström, N. O., Tran, T., Guo, J., Rutledge, E., Parvez, R. K., Thornton, M. E., Grubbs, B., McMahon, J. A. and McMahon, A. P. (2018). Conserved and divergent molecular and anatomic features of human and mouse nephron patterning. *J. Am. Soc. Nephrol.* **29**, 825-840.
- Little, M. H. and McMahon, A. P. (2012). Mammalian kidney development: principles, progress, and projections. *Cold Spring Harbor Perspect. Biol.* **4**, a008300.
- Lu, W., Shen, X., Pavlova, A., Lakkis, M., Ward, C. J., Pritchard, L., Harris, P. C., Genest, D. R., Perez-Atayde, A. R. and Zhou, J. (2001). Comparison of Pkd1-targeted mutants reveals that loss of polycystin-1 causes cystogenesis and bone defects. *Hum. Mol. Genet.* **10**, 2385-2396.
- Lyons, J. P., Miller, R. K., Zhou, X., Weidinger, G., Deroo, T., Denayer, T., Park, J.-I., Ji, H., Hong, J. Y., Li, A. et al. (2009). Requirement of Wnt/ β -catenin signaling in pronephric kidney development. *Mech. Dev.* **126**, 142-159.
- Maguire, R. J., Isaacs, H. V. and Pownall, M. E. (2012). Early transcriptional targets of MyoD link myogenesis and somitogenesis. *Dev. Biol.* **371**, 256-268.
- Matsumura, Y., Uchida, S., Kondo, Y., Miyazaki, H., Ko, S. B., Hayama, A., Morimoto, T., Liu, W., Arisawa, M., Sasaki, S. et al. (1999). Overt nephrogenic diabetes insipidus in mice lacking the CLC-K1 chloride channel. *Nat. Genet.* **21**, 95-98.
- Miner, J. H., Morello, R., Andrews, K. L., Li, C., Antignac, C., Shaw, A. S. and Lee, B. (2002). Transcriptional induction of slit diaphragm genes by Lmx1b is required in podocyte differentiation. *J. Clin. Investig.* **109**, 1065-1072.
- Møbjerg, N., Larsen, E. H. and Jespersen, Å. (2000). Morphology of the kidney in larvae of *Bufo viridis* (amphibia, anura, bufonidae). *J. Morphol.* **245**, 177-195.
- Moody, S. A. (1987). Fates of the blastomeres of the 32-cell-stage *Xenopus* embryo. *Dev. Biol.* **122**, 300-319.
- Murugan, S., Shan, J., Kühl, S. J., Tata, A., Pietilä, I., Kühl, M. and Vainio, S. J. (2012). WT1 and Sox11 regulate synergistically the promoter of the Wnt4 gene that encodes a critical signal for nephrogenesis. *Exp. Cell Res.* **318**, 1134-1145.
- Nagamani, S. C. S., Erez, A., Shen, J., Li, C., Roeder, E., Cox, S., Karaviti, L., Pearson, M., Kang, S. H. L., Sahoo, T. et al. (2010). Clinical spectrum associated with recurrent genomic rearrangements in chromosome 17q12. *Eur. J. Hum. Genet.* **18**, 278-284.
- Nakano, T., Niimura, F., Hohenfellner, K., Miyakita, E. and Ichikawa, I. (2003). Screening for mutations in BMP4 and FOXC1 genes in congenital anomalies of the kidney and urinary tract in humans. *Tokai J. Exp. Clin. Med.* **28**, 121-126.
- Naylor, R. W. and Jones, E. A. (2009). Notch activates Wnt-4 signalling to control medio-lateral patterning of the pronephros. *Development* **136**, 3585-3595.
- Neugebauer, J. M., Kwon, S., Kim, H.-S., Donley, N., Tilak, A., Sopory, S. and Christian, J. L. (2015). The prodomain of BMP4 is necessary and sufficient to generate stable BMP4/7 heterodimers with enhanced bioactivity in vivo. *Proc. Natl. Acad. Sci. USA* **112**, E2307-E2316.
- Nishinakamura, R. (2003). Kidney development conserved over species: Essential roles of Sall1. *Semin. Cell Dev. Biol.* **14**, 241-247.
- Ochi, H., Tamai, T., Nagano, H., Kawagushi, A., Suduo, N. and Ogino, H. (2012). Evolution of a tissue-specific silencer underlies divergence in the expression of pax2 and pax8 paralogues. *Nat. Commun.* **22**, 848.
- Olblich, H., Fliegau, M., Hoefele, J., Kispert, A., Otto, E., Volz, A., Wolf, M. T., Sasmaz, G., Trauer, U., Reinhardt, R. et al. (2003). Mutations in a novel gene, NPHP3, cause adolescent nephronophthisis, tapeto-retinal degeneration and hepatic fibrosis. *Nat. Genet.* **34**, 455-459.
- Prato, A. P., Musso, M., Ceccherini, I., Mattioli, G., Giunta, C., Ghiggeri, G. M. and Jasonni, V. (2009). Hirschsprung disease and congenital anomalies of the kidney and urinary tract (CAKUT): A novel syndromic association. *Medicine* **88**, 83-90.
- Quassinti, L., Maccari, E., Murri, O. and Bramucci, M. (2007). Comparison of ACE activity in amphibian tissues: *Rana esculenta* and *Xenopus laevis*. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **146**, 119-123.
- Raciti, D., Reggiani, L., Geffers, L., Jiang, Q., Bacchion, F., Subrizi, A. E., Clements, D., Tindal, C., Davidson, D. R., Kaissling, B. et al. (2008). Organization of the pronephric kidney revealed by large-scale gene expression mapping. *Genome Biol.* **9**, R84.
- Rankin, S. A., Zorn, A. M. and Buchholz, D. R. (2011). New doxycycline-inducible transgenic lines in *Xenopus*. *Dev. Dyn.* **240**, 1467-1474.
- Reggiani, L., Raciti, D., Airik, R., Kispert, A. and Brändli, A. W. (2007). The prepattern transcription factor *lrx3* directs nephron segment identity. *Genes Dev.* **21**, 2358-2370.
- Romaker, D., Kumar, V., Cerqueira, D. M., Cox, R. M. and Wessely, O. (2014). MicroRNAs are critical regulators of tuberous sclerosis complex and mTORC1 activity in the size control of the *Xenopus* kidney. *Proc. Natl. Acad. Sci. USA* **111**, 6335-6340.
- Roose, M., Sauer, K., Turan, G., Solomentsew, N., Werdien, D., Pramanik, K., Senkel, S., Ryffel, G. U. and Waldner, C. (2009). Heat-shock inducible cre strains to study organogenesis in transgenic *Xenopus laevis*. *Transgenic Res.* **18**, 595-605.
- Rosselot, C., Spraggon, L., Chia, I., Baturina, E., Riccio, P., Lu, B., Niederreither, K., Dolle, P., Duester, G., Chambon, P. et al. (2010). Non-cell-autonomous retinoid signaling is crucial for renal development. *Development* **137**, 283-292.
- Ruf, R. G., Xu, P.-X., Silvius, D., Otto, E. A., Beekmann, F., Muerb, U. T., Kumar, S., Neuhaus, T. J., Kemper, M. J., Raymond, R. M. et al. (2004). SIX1 mutations cause branchio-oto-renal syndrome by disruption of EYA1-SIX1-DNA complexes. *Proc. Natl. Acad. Sci. USA* **101**, 8090-8095.
- Sanna-Cherchi, S., Westland, R., Ghiggeri, G. M. and Gharavi, A. G. (2018). Genetic basis of human congenital anomalies of the kidney and urinary tract. *J. Clin. Investig.* **128**, 4-15.
- Sauer, K., Kahnert, S., Roose, M., Gull, M., Brandli, A. W., Ryffel, G. U. and Waldner, C. (2012). Heat-shock mediated overexpression of HNF1 β mutations has differential effects on gene expression in the *Xenopus* pronephric kidney. *PLoS ONE* **7**, e33522.
- Saulnier, D. M. E., Ghanbari, H. and Brändli, A. W. (2002). Essential function of Wnt-4 for Tubulogenesis in the *Xenopus* Pronephric Kidney. *Dev. Biol.* **248**, 13-28.
- Schmitt, S. M., Gull, M. and Brändli, A. W. (2014). Engineering *Xenopus* embryos for phenotypic drug discovery screening. *Adv. Drug Delivery. Rev.*, 225-246.
- Seys, E., Andriani, O., Keck, M., Mansour-Hendili, L., Courand, P.-Y., Simian, C., Deschenes, G., Kwon, T., Bertholet-Thomas, A., Bobrie, G. et al. (2017). Clinical and genetic spectrum of Bartter syndrome type 3. *J. Am. Soc. Nephrol.* **28**, 2540-2552.
- Shawlot, W. and Behringer, R. R. (1995). Requirement for *lim1* in head-organizer function. *Nature* **374**, 425-430.
- Shi, Z., Wang, F., Cui, Y., Liu, Z., Guo, X., Zhang, Y., Deng, Y., Zhao, H. and Chen, Y. (2015). Heritable CRISPR/Cas9-mediated targeted integration in *Xenopus* tropicalis. *FASEB J.* **29**, 4914-4923.
- Short, K. M., Combes, A. N., Lefevre, J., Ju, A. L., Georgas, K. M., Lamberton, T., Cairncross, O., Rumballe, B. A., McMahon, A. P., Hamilton, N. A. et al. (2014). Global quantification of tissue dynamics in the developing mouse kidney. *Dev. Cell* **29**, 188-202.
- Simon, D. B., Bindra, R. S., Mansfield, T. A., NelsonWilliams, C., Mendonca, E., Stone, R., Schurman, S., Nayir, A., Alpay, H., Bakkaloglu, A. et al. (1997). Mutations in the chloride channel gene, *CLCNKB*, cause Bartter's syndrome type III. *Nat. Genet.* **17**, 171-178.
- Sinner, D., Kirilenko, P., Rankin, S., Wei, E., Howard, L., Kofron, M., Heasman, J., Woodland, H. R. and Zorn, A. M. (2006). Global analysis of the transcriptional network controlling *Xenopus* endoderm formation. *Development* **133**, 1955-1966.

- Skinner, M. A., Safford, S. D., Reeves, J. G., Jackson, M. E. and Freemerman, A. J. (2008). Renal aplasia in humans is associated with RET mutations. *Am. J. Hum. Genet.* **82**, 344-351.
- Sorokin, L., Klein, G., Mugrauer, G., Fecker, L., Ekblom, M. and Ekblom, P. (1992). Development of kidney epithelial cells. In *Epithelial Organization and Development* (ed T.P. Fleming), pp. 163-187. Springer, Dordrecht.
- Srivastava, S., Molinari, E., Raman, S. and Sayer, J. A. (2018). Many genes—one disease? Genetics of Nephronophthisis (NPHP) and NPHP-associated disorders. *Front. Pediatr.* **5**, 1-15.
- Stokman, M., Lilien, M. and Knoers, N. (2016). Nephronophthisis. *Gene Reviews*.
- Suspitsin, E. N. and Imyanitov, E. N. (2016). Bardet-biedl syndrome. *Mol. Syndromol.* **7**, 62-71.
- Suzuki, N., Hirano, K., Ogino, H. and Ochi, H. (2015). Identification of distal enhancers for Six2 expression in pronephros. *Int. J. Dev. Biol.* **59**, 241-246.
- Suzuki, N., Hirano, K., Ogino, H. and Ochi, H. (2019). Arid3a regulates nephric tubule regeneration via evolutionarily conserved regeneration signal-response enhancers. *eLife* **8**, e43186.
- Taskiran, E. Z., Korkmaz, E., Gucer, S., Kosukcu, C., Kaymaz, F., Koyunlar, C., Bryda, E. C., Chaki, M., Lu, D., Vadnagara, K. et al. (2014). Mutations in ANKS6 Cause a Nephronophthisis-Like Phenotype with ESRD. *J. Am. Soc. Nephrol.* **25**, 1653-1661.
- Tena, J. J., Neto, A., de la Calle-Mustienes, E., Bras-Pereira, C., Casares, F. and Gómez-Skarmeta, J. L. (2007). Odd-skipped genes encode repressors that control kidney development. *Dev. Biol.* **301**, 518-531.
- Tran, U., Pickney, L. M., Özpolat, B. D. and Wessely, O. (2007). Xenopus Bicaudal-C is required for the differentiation of the amphibian pronephros. *Dev. Biol.* **307**, 152-164.
- Tran, U., Zakin, L., Schweickert, A., Agrawal, R., Doger, R., Blum, M., De Robertis, E. M. and Wessely, O. (2010). The RNA-binding protein bicaudal C regulates polycystin 2 in the kidney by antagonizing miR-17 activity. *Development* **137**, 1107-1116.
- Urban, A. E., Zhou, X., Ungos, J. M., Raible, D. W., Altmann, C. R. and Vize, P. D. (2006). FGF is essential for both condensation and mesenchymal-epithelial transition stages of pronephric kidney tubule development. *Dev. Biol.* **297**, 103-117.
- Van Esch, H., Groenen, P., Nesbit, M. A., Schuffenhauer, S., Lichtner, P., Vanderlinden, G., Harding, B., Beetz, R., Bilous, R. W., Holdaway, I. et al. (2000). GATA3 haplo-insufficiency causes human HDR syndrome. *Nature* **406**, 419-422.
- Verdeguer, F., Le Corre, S., Fischer, E., Callens, C., Garbay, S., Doyen, A., Igarashi, P., Terzi, F. and Pontoglio, M. (2010). A mitotic transcriptional switch in polycystic kidney disease. *Nat. Med.* **16**, 106-110.
- Vivante, A. and Hildebrandt, F. (2016). Genetics of congenital anomalies of the kidneys and urinary tract. In *Congenital Anomalies of the Kidney and Urinary Tract: Clinical Implications in Children* (eds Barakat A., Rushton H.), pp 303-322. Springer, Cham.
- Vivante, A., Mark-Danieli, M., Davidovits, M., Harari-Steinberg, O., Omer, D., Gnatek, Y., Cleper, R., Landau, D., Kovalski, Y., Weissman, I. et al. (2013). Renal hypodysplasia associates with a Wnt4 variant that causes aberrant canonical Wnt signaling. *J. Am. Soc. Nephrol.* **24**, 550-558.
- Vivante, A., Kohl, S., Hwang, D.-Y., Dworschak, G. C. and Hildebrandt, F. (2014). Single-gene causes of congenital anomalies of the kidney and urinary tract (CAKUT) in humans. *Pediatr. Nephrol.* **29**, 695-704.
- Vivante, A., Mann, N., Yonath, H., Weiss, A.-C., Getwan, M., Kaminski, M. M., Bohnenpoll, T., Teyssier, C., Chen, J., Shril, S. et al. (2017). A dominant mutation in nuclear receptor interacting protein 1 causes urinary tract malformations via dysregulation of retinoic acid signaling. *J. Am. Soc. Nephrol.* **28**, 2364-2376.
- Vize, P. D. (2003). The chloride conductance channel CIC-K is a specific marker for the Xenopus pronephric distal tubule and duct. *Gene Exp. Patterns* **3**, 347-350.
- Vize, P. D., Carroll, T. J. and Wallingford, J. B. (2003). Induction, development, and physiology of the pronephric tubules. In *The Kidney* (eds P. D. Vize, A. S. Woolf and J. B.L. Bard), pp. 19-50. Elsevier Academic Press.
- Wang, S., Krinks, M., Kleinwaks, L. and Moos, M. (1997). A novel Xenopus homologue of bone morphogenetic protein-7 (BMP-7). *Genes Funct.* **1**, 259-271.
- Waters, A. M. and Beales, P. L. (2011). Ciliopathies: an expanding disease spectrum. *Pediatr. Nephrol.* **26**, 1039-1056.
- Weber, S., Taylor, J. C., Winyard, P., Baker, K. F., Sullivan-Brown, J., Schild, R., Knuppel, T., Zurowska, A. M., Caldas-Alfonso, A., Litwin, M. et al. (2008). SIX2 and BMP4 mutations associate with anomalous kidney development. *J. Am. Soc. Nephrol.* **19**, 891-903.
- Wessely, O. and Tran, U. (2011). Xenopus pronephros development—past, present, and future. *Pediatr. Nephrol.* **26**, 1545-1551.
- White, J. T., Zhang, B., Cerqueira, D. M., Tran, U. and Wessely, O. (2010). Notch signaling, wt1 and foxc2 are key regulators of the podocyte gene regulatory network in Xenopus. *Development* **137**, 1863-1873.
- Wing, M. R., Devaney, J. M., Joffe, M. M., Xie, D., Feldman, H. I., Dominic, E. A., Guzman, N. J., Ramezani, A., Susztak, K., Herman, J. G. et al. (2014). DNA methylation profile associated with rapid decline in kidney function: Findings from the CRIC Study. *Nephrol. Dial. Transplant.* **29**, 864-872.
- Wolf, M. T. F. and Hildebrandt, F. (2011). Nephronophthisis. *Pediatric Nephrol.* **26**, 181-194.
- Xue, Y., Chen, Y., Ayub, Q., Huang, N., Ball, E. V., Mort, M., Phillips, A. D., Shaw, K., Stenson, P. D., Cooper, D. N. et al. (2012). Deleterious- and disease-allele prevalence in healthy individuals: Insights from current predictions, mutation databases, and population-scale resequencing. *Am. J. Hum. Genet.* **91**, 1022-1032.
- Yan, B., Neilson, K. M., Ranganathan, R., Maynard, T., Streit, A. and Moody, S. A. (2015). Microarray identification of novel genes downstream of Six1, a critical factor in cranial placode, somite, and kidney development. *Dev. Dyn.* **244**, 181-210.
- Yildirim-Toruner, C., Subramanian, K., El Manjra, L., Chen, E., Goldstein, S. and Vitale, E. (2004). A novel frameshift mutation of FOXC2 gene in a family with hereditary lymphedema-distichiasis syndrome associated with renal disease and diabetes mellitus. *Am. J. Med. Genet. A* **131**, 281-286.
- Yosypiv, I. V. (2012). Congenital anomalies of the kidney and urinary tract: a genetic disorder? *Int. J. Nephrol.* **2012**, 909083.
- Zalli, D., Bayliss, R. and Fry, A. M. (2012). The Nek8 protein kinase, mutated in the human cystic kidney disease nephronophthisis, is both activated and degraded during ciliogenesis. *Hum. Mol. Genet.* **21**, 1155-1171.
- Zhou, X. and Vize, P. D. (2004). Proximo-distal specialization of epithelial transport processes within the Xenopus pronephric kidney tubules. *Dev. Biol.* **271**, 322-338.
- Zhou, X. and Vize, P. D. (2005). Pronephric regulation of acid-base balance; coexpression of carbonic anhydrase type 2 and sodium-bicarbonate cotransporter-1 in the late distal segment. *Dev. Dyn.* **233**, 142-144.
- Zhuo, X., Haeri, M., Solessio, E. and Knox, B. E. (2013). An inducible expression system to measure rhodopsin transport in transgenic Xenopus rod outer segments. *PLoS ONE* **8**, e82629.