

Haploinsufficiency of the murine polycomb gene *Suz12* results in diverse malformations of the brain and neural tube

Xavier Miró^{1,*}, Xunlei Zhou², Susann Boretius³, Thomas Michaelis³, Christian Kubisch^{4,5,6}, Gonzalo Alvarez-Bolado² and Peter Gruss¹

SUMMARY

Polycomb proteins are epigenetic regulators of gene expression. Human central nervous system (CNS) malformations are congenital defects of the brain and spinal cord. One example of a human CNS malformation is Chiari malformation (CM), which presents as abnormal brainstem growth and cerebellar herniation, sometimes accompanied by spina bifida and cortical defects; it can occur in families. Clinically, CM ranges from an asymptomatic condition to one with incapacitating or lethal symptoms, including neural tube defects and hydrocephalus. However, no genes that are causally involved in any manifestation of CM or similar malformations have been identified. Here, we show that a pathway that involves *Zac1* (also known as *Plag1* or *Lot1*) and controls neuronal proliferation is altered in mice that are heterozygous for the polycomb gene *Suz12*, resulting in a phenotype that overlaps with some clinical manifestations of the CM spectrum. *Suz12* heterozygotes show cerebellar herniation and an enlarged brainstem, accompanied by occipital cortical alterations and spina bifida. Downward displacement of the cerebellum causes hydrocephalus in the most severely impaired cases. Although the involvement of polycomb genes in human disease is starting to be recognized, this is the first demonstration of their role in nervous system malformations. Our work strongly suggests that brain malformations such as CM can result from altered epigenetic regulation of genes involved in cell proliferation in the brain.

RESULTS AND DISCUSSION

Polycomb proteins work in multimeric complexes termed polycomb repressive complexes (PRCs) to regulate the expression of numerous developmental genes that control cell proliferation and stem cell identity, as well as genomic imprinting and X-chromosome inactivation, and are involved, for example, in cancer development (Lee et al., 2006; Sparmann and van Lohuizen 2006; Schuettengruber et al., 2007). *Suz12*, *Eed* and *Ezh2* are polycomb proteins that combine in PRC2 to silence genes (Cao et al., 2002). In the mouse, *Suz12* expression starts with a peak during gastrulation [embryonic day (E)5.5 to E8.5]. This early expression is essential for survival, since *Suz12* knockout mice die at gastrulation (as do *Eed*^{-/-} and *Ezh2*^{-/-} embryos) (Faust et al., 1998; O'Carroll et al., 2001; Pasini et al., 2004). We detected a second period of intense expression in several organs, including the brain, where expression continues unabated through development and into the adult (Fig. 1A,B). At midgestation, *Suz12* was expressed in a posterior-to-anterior gradient in the primordium of the cerebral cortex, as well as in the tectum, hindbrain, eye primordium, somites, first branchial arch, nasal cavity, kidney, lung, liver and pancreas (Fig. 1C-K,O,P). Later, *Suz12* transcripts appeared in the

cerebellum and hippocampus (Fig. 1L-N). We also found transcripts of *Ezh2* and *Eed* colocalizing with *Suz12* in the developing cortex, tectum, nasal epithelia, liver and kidney (Fig. 1Q,R), suggesting a role for PRC2 in the development of these organs.

The role of the late peak of expression has not yet been addressed. However, *Drosophila* mutants that are heterozygous for *Su(z)12* show major morphological alterations, including homeotic transformations (Birve et al., 2001). Therefore, in order to investigate the role of *Suz12* in mouse development beyond gastrulation, we decided to generate a *Suz12*-deficient mouse line (Fig. 2A,B). The homozygous knockout animals showed an early lethal phenotype, as reported previously (Pasini et al., 2004). The heterozygotes, however, survived and reproduced, although they showed a range of central nervous system (CNS) abnormalities, which were not gender related (Fig. 2C-D"). The consistent increases and decreases in size of specific brain regions were particularly intriguing. In contrast to the anterior cortex (A-CTX) and other normal-sized telencephalic regions, such as the caudate-putamen and hippocampus, the posterior cortex (P-CTX) was smaller in *Suz12*^{+/-} mice than in wild-type animals (cortical heterometry) (Fig. 2C-G; supplementary material Fig. S1). This cortical heterometry correlated well with the caudo-rostral gradient of *Suz12* expression (Fig. 1D,E,P).

We then examined the expression of two marker genes that are expressed specifically in the P-CTX during development, *Id2* and *COUP-TF1* (also known as *Nr2f1*) (Armentano et al., 2006; Lasorella et al., 2006). Both of these genes showed shortened domains of expression in the heterozygous mutants, indicating that the cortical reduction in size affects specifically the P-CTX (Fig. 2H-K).

By contrast, there was a consistent increase in the size of the midbrain tectum of heterozygous animals, where both the superior

¹Max Planck Institute of Biophysical Chemistry, Department of Molecular Cell Biology, Göttingen D-37077, Germany

²Max Planck Institute of Biophysical Chemistry, Department of Genes and Behavior, Brain Development Group, Göttingen D-37077, Germany

³Biomedizinische NMR Forschungs GmbH, Max Planck Institute of Biophysical Chemistry, Göttingen D-37077, Germany

⁴Institute of Human Genetics, ⁵Institute for Genetics and ⁶Center of Molecular Medicine Cologne (CMCC), University of Cologne, Cologne D-50931, Germany

*Author for correspondence (e-mail: xmiro@gwdg.de)




Fig. 1. Expression of *Suz12* in mouse development.

(A) Detection of *Suz12* expression through mouse embryo development by northern blotting. Stages E4.5-E6.5 include extra-embryonic tissues and maternal uterus; stages E7.5-E9.5 include embryo and extra-embryonic tissues. There are two peaks of *Suz12* expression, including one at the very early stages of development. (B) The *Suz12* transcript remains expressed in mouse brain up to older ages, as shown by northern blotting. (C-K) Expression of *Suz12* in the mouse embryo. β -galactosidase (β -gal) staining on whole-mount *Suz12*^{+/-} mouse embryos at E10.5 (F) and E11.5 (G) shows strong expression in the developing first branchial arch (br) and liver (li). In situ hybridization (ISH) of *Suz12* on coronal (C) and sagittal (D,E,H,J,P) sections at E12.5 (C,D,H,J) and at E14.5 (E,P) correlates with the β -gal staining on whole-mount *Suz12*^{+/-} mouse embryos at E12.5 (I) and E18.5 (K), and with β -gal ISH of *Suz12*^{+/-} mouse embryos at E14.5 (O). The *Suz12* transcript shows a posterior (p)-to-anterior (a) gradient of expression in the primordium of the cerebral cortex (D,E,I,O,P), with no expression in the cortical hem (hem) (C-E), and expression in the tectal neuroepithelium (tc) (J,K), eye primordium (H,I), nasal epithelium (ne), kidney (ki), lung (lu) and pancreas (pa) (O,P). (L-N) β -gal ISH (L) and X-gal reaction (M,N) of *Suz12*^{+/-} mouse embryos at P9 reveals *Suz12* expression in the external granular layer (egl) of the cerebellum, as well as in the Purkinje cell layer (pcl) (L), which represents the only expression in the adult cerebellum (M). *Suz12* is also expressed in the hippocampus, with enrichment in the adult CA2 (N). (Q,R) Transcripts of *Ezh2* (Q) and *Eed* (R), two other PRC2 partners, show an expression pattern similar to *Suz12*. Bars, 250 μ m (C,D,I-N); 100 μ m (E,H); 500 μ m (F,G); 1 mm (O-R).

and inferior colliculi (SC and IC, respectively) were enlarged significantly (Fig. 2E-G; supplementary material Fig. S1). These phenotypical traits co-segregated so that the individuals with the largest reduction in the size of the P-CTX showed, at the same time, the largest increase in IC and SC surface area. Over 60% of the mutants measured had values that were outside the wild-type range (supplementary material Fig. S1). The overall changes in the P-CTX, IC and SC were statistically significant (Fig. 2G).

The enlargement of the IC and SC produced a 'beaked tectum' (Fig. 3A,B), which, in the most pronounced cases, caused dramatic cerebellar herniation through the foramen magnum and a secondary hydrocephalus (Fig. 3C,D). Magnetic resonance imaging (MRI) of living *Suz12* heterozygotes (Fig. 3E-H) showed herniation

of the cerebellar paraflocculus (corresponding to the human cerebellar tonsils) into the foramen magnum (yellow arrowhead in Fig. 3E,G), with a crowded posterior fossa (white arrowheads in Fig. 3E,G).

Next, we asked what mechanism could produce the alterations in size that we found in the *Suz12* heterozygote brain. The decrease in cortex size or increase in tectum size could be attributed to a dysregulation of cell proliferation. Indeed, bromodeoxyuridine (BrdU)-labeling of brains in heterozygous animals showed strongly reduced proliferation in the P-CTX (Fig. 3I) (apoptosis, as detected by TUNEL, was unchanged; data not shown) and increased proliferation in the tectum (Fig. 3I). Since the role of the PRC is to maintain transcriptional repression, the defects observed in *Suz12*




Fig. 2. *Suz12*^{+/-} mice show a reduced occipital cortex and enlarged tectum. (A,B) Inactivation of *Suz12*. (A) Schematic representation of wild-type *Suz12*^{+/+} (top) and mutated *Suz12*^{-/-} (bottom) alleles. Exons are represented by black boxes. The sizes of the restriction fragments that were expected to be recognized by the Southern probe (indicated) are shown. (B) Southern blot. (C-D'') *Suz12*^{+/-} mice show different degrees of brain reduction. A comparison between the brains of 46-day-old wild-type (C,D) and *Suz12*^{+/-} (C',C'',D',D'') mice shows different degrees of reduction in the size of the P-CTX of the heterozygote (arrowheads). D-D'' show Nissl-stained sections of C-C'' brains. (E-G) Histological measurements show a specific reduction of the P-CTX and enlargement of the tectum in the newborn *Suz12*^{-/-} brain (discordant asymmetry). (E,F) We subdivided the cortex into two parts, anterior (A-CTX) and posterior (P-CTX), and measured them, as well as the inferior and superior colliculus (IC and SC, respectively), hippocampus (HIPP) and caudate-putamen (CP). (G) Statistical analysis showed that *Suz12*^{+/-} brains had a reduced P-CTX, as well as an enlarged IC and SC (tectum), with no differences in the other regions. (H-K) Expression of the posterior cortical developmental markers, *Id2* (H,I) and *COUP-TFI* (J,K), showed shortened domains in the *Suz12*^{+/-} cortex (I,K) compared with the wild-type cortex (H,J). Arrowheads indicate the anterior boundary of the expression domain; OB, olfactory bulb (for reference). The values in G represent the mean ± s.d. of measurements from seven wild-type and 13 mutant brains. Bars, 1 mm (C-E); 250 μm (H-K).

heterozygous mice could have resulted from the inappropriate expression of genes involved in neural stem cell proliferation. Some targets of SUZ12 in human embryonic stem cells have recently been mapped by chromatin immunoprecipitation on a genome-wide scale (Lee et al., 2006). One of those targets, *ZAC* (also known as *PLAGL1* or *LOT1*), encodes a transcription factor that regulates

apoptosis and cell cycle arrest in the neuroepithelium (Varrault et al., 2006). Intriguingly, embryonic and postnatal expression of the murine ortholog of *ZAC*, *Zac1*, colocalized strongly with that of *Suz12* (compare Fig. 3J with Fig. 1P) (Valente and Auladell, 2001). To verify this point, we labeled *Suz12* and *Zac1* with specific antibodies on sections of E14.5 wild-type brain. The results show




Fig. 3. Beaked tectum, cerebellar herniation and hydrocephalus, as well as altered *Zac1* expression, in *Suz12*^{+/-} mice. (A,B) An enlarged beaked tectum (tc) in *Suz12*^{+/-} mice (B) results in herniation of the cerebellum when compared with wild-type littermates (A). In the most dramatic herniations, hydrocephalus (HYD) was observed (C,D), including severe cerebellar (CB) abnormalities (D). (E-H) MRI studies of 42-day-old mice. The comparison between sagittal views of T2-weighted MRIs of wild-type (E,F) and *Suz12*^{+/-} (G,H) mice shows how the ventral paraflocculus (the structure that corresponds to the cerebellar tonsils in humans) goes into the foramen magnum in *Suz12*^{+/-} mice as a consequence of cerebellar herniation (yellow arrowhead) (F,G). At the same time, the *Suz12*^{+/-} mice have a crowded posterior fossa, with the volumes of both the cisterna cerebellomedullaris and pontis reduced (white arrowheads) (F,G). (I) BrdU labeling showed strongly reduced proliferation, specifically in the P-CTX (but not the A-CTX), as well as increased proliferation in the tectum (TC). (J-P) *Zac1* is specifically upregulated in areas of altered proliferation in the *Suz12*^{+/-} brain. *Zac1* shows an expression pattern that is very similar to *Suz12*, as observed by ISH on *Suz12*^{+/-} mice at E14.5 (J). *Zac1* (red in K) and *Suz12* (green in K) colocalize in the cortical neuroepithelium (ctx) but not in the hem (K). *Zac1* is upregulated in the *Suz12*^{+/-} cortex (CTX) (L-N) and tectal central gray neuroepithelium (tcn) (O,P), as revealed by ISH (J,L,M,O,P) and real-time PCR (N). In addition, real-time PCR revealed increasing expression of *Zac1* in *Suz12*^{+/-} E13.5 embryonic fibroblasts (MEF) (N). (Q) *SUZ12* knockdown in human osteosarcoma U2Os cells resulted in upregulation of ZAC. Transfection with specific siRNA dramatically reduced *SUZ12* expression, as confirmed by western blotting and real-time PCR (Q), resulting in upregulation of ZAC at both the protein and mRNA levels (Q). (R,S) Detection of P-Creb by immunohistochemistry at E14.5 revealed an increase in the *Suz12*^{+/-} P-CTX (arrowheads). The values in I represent the mean±s.d. of measurements on three brains; the values in N,Q represent the mean±s.d. of three experiments. Bars, 500 μm (A-D,F,G); 1 mm (E,H,J); 100 μm (K,R,S); 250 μm (L,M,O,P).

that both proteins are colocalized in the cortical neuroepithelium (ctx in Fig. 3K). The hem region lacks the *Suz12* protein (Fig. 3K), in agreement with our mRNA data (Fig. 1E).

Zac1 was upregulated in the *Suz12* heterozygous cortex (by real-time PCR and by ISH) (Fig. 3L-N) and tectum (Fig. 3O,P). Cultured fibroblasts from heterozygous embryos also showed *Zac1* upregulation (Fig. 3N). To verify a link between the downregulation of *Suz12* and the increase in *Zac1* expression, we used small interfering RNA (siRNA) to knockdown *SUZ12* in a cell line that is commonly used for silencing analysis of the polycomb genes (Aoto et al., 2008), the human osteosarcoma U2Os line (U2Os-*Suz12* RNAi cells). Knocking down *SUZ12* in these cells resulted in upregulation of ZAC at the protein and mRNA levels (Fig. 3Q). *Zac1* exerts its effects by inducing the expression of *Pac1* (also known as *Adcyap1r1*) (Rodriguez-Henche et al., 2002), a receptor present in the neuroepithelium. Injection of the *Pac1* ligand

pituitary adenylate cyclase-activating polypeptide (PACAP) into the developing rat cortex decreases proliferation (without increasing apoptosis) and enhances the phosphorylation of the cAMP-responsive element-binding protein (CREB) (Suh et al., 2001). In agreement with this, we detected increased expression of phosphorylated (P)-Creb in the P-CTX of heterozygous mice at E14.5 (Fig. 3R,S), supporting our conclusion that *Suz12* acts in normal cortical development, at least in part, by regulating the *Zac1-Pac1-Creb1* pathway.

Additional phenotypical alterations that are found in the heterozygous mice (see below) could be caused through a dysregulation of the expression of different genes that are also normally silenced by *Suz12*. Indeed, the polycomb proteins, including *Suz12*, are known to regulate the spatial patterns of homeotic box (Hox) gene expression in *Drosophila* (Cao and Zhang, 2004; Schuettengruber et al., 2007) and, in vertebrates, a Hox gene

code controls the orderly differentiation of vertebrae (Kessel and Gruss, 1990). The most seriously affected of our heterozygous mice showed a protuberated back (white arrowhead in Fig. 4A) and varying degrees of spina bifida (Fig. 4E-G). These mutants subsequently developed paraplegia (yellow arrowhead in Fig. 4A). *Suz12* is indeed expressed in the early neural tube (Fig. 4B-D). Diastematomyelia, a vertical splitting of the spinal cord by a bony, fibrous or cartilaginous septum (Parmar et al., 2003), was also observed in lumbar L3-L4 vertebrae (Fig. 4H), the development of which is controlled by *Hoxd9* (Burke et al., 1995). Consistently, it has been shown that *SUZ12* binds to the *HOXD9* promoter region in human embryonic stem cells (Lee et al., 2006). Finally, some of the *Suz12* heterozygotes showed hippocampal morphological alterations (Fig. 4I,J), as well as partial agenesis of the corpus callosum (arrows in Fig. 4K,L). We also observed a loss of gray matter in a gradient matching the posterior-to-anterior gradient of *Suz12* expression (see above), together with layer disorganization in the P-CTX (Fig. 4K,L).

The morphological CNS alterations show incomplete penetrance and variable expressivity in the heterozygous mice, and are also found in varying degrees of severity in human patients with neural tube defects and congenital brain malformations (Adeloye, 1976). These severely disabling and, as a group, rather common pathologies represent a continuum of disorders with multifactorial etiology. One of the sub-phenotypes is Chiari malformation (CM), which typically shows herniation of the cerebellar tonsils through

the foramen magnum with secondary hydrocephalus. Additional symptoms may include diastematomyelia (Parmar et al., 2003); secondary paralysis of the legs (Stevenson, 2004); abnormal gyral pattern and dysplasia in occipital regions (Stevenson, 2004); and partial agenesis of the corpus callosum. Thus, this syndrome shows partial overlap with symptoms in some of the haploinsufficient *Suz12* mice, making a causative role of *SUZ12* in human CNS malformations an attractive hypothesis. However, it has to be noted that none of these clinically defined disease entities follows a monogenic mode of inheritance in humans, making it difficult to obtain definite proof that *SUZ12* or other genes of the polycomb complex have a causative genetic involvement in disease pathophysiology. Moreover, the Chiari-like phenotype in our mice seems to be the result of the gross enlargement of the tectum, which has not been described as a characteristic primary alteration in CM or other clinically defined human CNS malformation syndromes, leaving the comparability of phenotypes unclear. Nevertheless, a direct or indirect involvement of polycomb-mediated epigenetic changes in CNS malformations still seems attractive, particularly because of the proven association between neural tube defects and folate-dependent methylation pathways (see overview in Kibar et al., 2007).

Additional evidence supporting a role for *SUZ12* in human CNS malformations may come from clinical and genetic studies on neurofibromatosis type 1 (NF1), an autosomal dominant phakomatosis affecting 1 in 3500 individuals, which is caused by a heterozygous mutation in the tumor suppressor gene *NF1* (Venturin et al., 2004). In addition to the characteristic tumor predisposition and cutaneous symptoms, some NF1 patients do also show severe mental retardation and brain malformations, including disturbances of cortical development (Balestri et al., 2003). The majority of NF1 cases are caused by point mutations in the *NF1* gene. However, 5-10% of cases show a larger genomic deletion (Venturin et al., 2004), comprising the *SUZ12* and around 12 additional genes, typically resulting in a more severe phenotype. This aggravated form of the disease is probably because of the haploinsufficiency of one or more of the additionally deleted genes, including *SUZ12*. Moreover, some NF1 patients have been found to show a CM (Tubbs et al., 2004), although a possible link to the type of underlying mutation has not been investigated in this study. By contrast, there is a small study that describes five independent NF1 patients with microdeletions, all of whom show severe developmental impairment and three of whom have structural brain anomalies (including one patient with a CM and hydrocephalus, and another with dysgenesis of the corpus callosum) (Korf et al., 1999). These studies would be compatible with the hypothesis that haploinsufficiency of *SUZ12* may be involved in the development of the more severe phenotype of NF1 microdeletion patients and possibly in the development of diverse brain malformations or, more generally, brain malfunction.

Epigenetic alterations are, on the one hand, essential for proper development but, on the other hand and in the case of dysregulation, they are increasingly being linked to a variety of human diseases (Feinberg, 2007). For example, the involvement of alterations in polycomb silencing in the pathogenesis of cancer is an active area of research (Tan et al., 2007). Moreover, our results are the first to indicate that a genetically induced alteration of epigenetic transcriptional control mechanisms can underlie




Fig. 4. Spina bifida, diastematomyelia and callosal alterations in *Suz12*^{+/-} mice. (A) In the most impaired cases, *Suz12*^{+/-} mice showed a protuberated back (white arrowhead) and paraplegia of the posterior legs (yellow arrowhead). (B-D) *Suz12* expression in the neural tube (NT) at E8.5 (B), E9.5 (C) and E10.5 (D). (E-H) Malformations observed in the vertebral column of *Suz12*^{+/-} mice. Diverse degrees of spina bifida (black arrows) were observed in heterozygous mice (F,G) in contrast to wild-type mice (E). In addition, diastematomyelia (H) was also found, caused by bony septa in lumbar L3-L4 vertebrae (white arrowheads). (I-L) Other alterations observed in the brain were deformations of the hippocampal cortex (I,J), partial agenesis of the corpus callosum (black arrows) (K,L), and loss of the gray matter in a gradient (K,L; see also Fig. 3C) that follows the posterior-to-anterior *Suz12* expression gradient (Fig. 1D,E), with disorganization of cortical layers in the P-CTX of 24-day-old *Suz12*^{+/-} mice compared with wild-type brains (K,L). Bars, 500 μ m (B-F,I-L); 1 mm (G); 250 μ m (H).

TRANSLATIONAL IMPACT

Clinical issue

Brain and neural tube defects are among the most common human congenital malformations. One prominent example is the Chiari malformation (CM), a developmental defect characterized by abnormal growth of the brainstem and cerebellar herniation. CMs can be asymptomatic or can cause a wide range of clinical conditions. Malformations may, for instance, cause severe neural tube defects and hydrocephalus, in turn leading to incapacitating or lethal conditions. CMs and most other congenital brain defects have a pronounced genetic basis and a complex inheritance pattern. The genetic causes of these defects, and the molecular interplay between known environmental risk factors (for example, folic acid deficiency) and susceptibility genes are currently ill defined.

Results

In this study, the authors identify two genetic factors that cause Chiari-like malformations in mice. One of these genes, *Suz12*, is a member of a polycomb complex, a group of proteins that act as epigenetic regulators of gene expression. In comparison to normal, wild-type mice, which have two copies of *Suz12*, mice with only one copy of *Suz12* have brain and neural tube defects. Some of the *Suz12* heterozygotes show cerebellar herniation and an enlarged brainstem accompanied by occipital cortical alterations and spina bifida, almost perfectly resembling CM as seen in some human patients. The most impaired mice have downward displacement of the cerebellum, which causes hydrocephalus. The authors also demonstrate that *Zac1* (also known as *Plagl1* or *Lot1*), a regulator of neuronal proliferation, is part of the same molecular pathway affected by *Suz12* deficiency.

Implications and future directions

Polycomb proteins are starting to be recognized as having a key role in human disease; this study demonstrates that polycomb protein complexes can play a role in nervous system malformations as well. This work strongly suggests that developmental brain abnormalities such as CMs can result from altered epigenetic regulation of genes involved in cell proliferation in the brain.

doi:10.1242/dmm.003665

congenital malformations of the brain and spinal cord, which represent a diverse group of rather common and severely disabling human disorders of the CNS.

METHODS

Suz12 mutant mice

Using a genomic clone containing *Suz12* from mouse library PAC-RPCI-21 (RPCIP711PP78, German Resource Center for Genomics), we targeted exon 12 with lacZ and the PGK-neomycin cassette (Fig. 2A), and then generated *Suz12* mutant mice as described previously (Nagy et al., 2003). Animals were handled according to European and German law. PCR primers for wild-type (0.35 kb) and mutant (0.30 kb) alleles: P1, 5'-GCCTGAAGAACGAGATCA-3'; P2, 5'-CCAGGTCATCTTGTGGAG-3'; P3, 5'-TGGAGCTGGAGT-TACCTG-3'. Northern blot probe: nucleotides 3163-3937 of GenBank ID BC064461 (*Suz12*).

Antibodies

Primary antibodies: anti-SUZ12 (Upstate), anti-BrdU (Dako), anti-phospho-CREB (Upstate), anti- β -tubulin (Santa Cruz), anti-Zac1 (Santa Cruz). Secondary antibodies: biotinylated universal anti-mouse/rabbit IgG (ABC) and HRP-conjugated anti-rabbit IgG (Amersham).

ISH probes

IMAGE consortium CloneID (Lennon et al., 1996) and GenBank Accession ID for ISH probes: *Suz12*, IRAKp961B0154Q2 (BC064461, nucleotides 3163-4448); *Ezh2*, IRAKp961I0810Q2 (BC016391, 1576-2595). For *Zac1*, *Eed*, *LacZ* and *COUP-TFI*, we amplified probes from E14.5 mouse brain cDNA, covering nucleotides: 1838-2636 of AK087432 (*Zac1*), 529-1531 of AK077664 (*Eed*), 324-897 of U89671 (*LacZ*), and 621-1413 of U07625 (*COUP-TFI*). For *Id2*, RIKEN clone 2810464O09 was used (Kawai et al., 2001) (<http://genome.gsc.riken.go.jp/>).

MRI

T2-weighted images [2D fast spin-echo (FSE), repetition time (TR)=7 seconds, effective echo time (TE)=42.7 milliseconds, 8 echos, 50 contiguous slices] were obtained in sagittal and axial planes, at a magnetic field strength of 9.4 Tesla (Bruker Biospin GmbH, Germany), at a spatial resolution of 100×100×250 μm^3 .

Other methods

Brain areas were measured with Scion Image software (Scion Corporation). The real-time PCR primers for *Suz12* were 5'-CTTCCCCTGCAAACGAAG-3' and 5'-CCCCTGAGACAC-TATCTG-3'. The RNAi *SUZ12*-specific sequence was as described (Pasini et al., 2004). For statistics, Student's *t*-test was used; $P \leq 0.05$ was considered significant.

ACKNOWLEDGEMENTS

This work was supported by the Max Planck Society.

COMPETING INTERESTS

The authors declare no competing financial interests.

AUTHOR CONTRIBUTIONS

X.M., G.A.-B. and P.G. conceived and designed the experiments; X.M. performed the experiments; X.Z. contributed in the generation of the *Suz12*-deficient mice; S.B. and T.M. performed the MRI assays; X.M. and G.A.-B. analyzed the data; X.M., G.A.-B. and C.K. wrote the paper.

SUPPLEMENTARY MATERIAL

Supplementary material for this article is available at <http://dmm.biologists.org/lookup/suppl/doi:10.1242/dmm.001602/-/DC1>

Received 19 September 2008; Accepted 20 March 2009.

REFERENCES

- Adeloye, A. (1976). Mesencephalic spur (beaking deformity of the tectum) in Arnold-Chiari malformation. *J. Neurosurg.* **45**, 315-320.
- Aoto, T. N., Saitoh, N., Sakamoto, Y., Watanabe, S. and Nakao, M. (2008). Polycomb group protein-associated chromatin is reproduced in post-mitotic G1 phase and is required for S phase progression. *J. Biol. Chem.* **283**, 18905-18915.
- Armentano, M., Filosa, A., Andolfi, G. and Studer, M. (2006). COUP-TFI is required for the formation of commissural projections in the forebrain by regulating axonal growth. *Development* **133**, 4151-4162.
- Balestri, P., Vivarelli, R., Grosso, S., Santori, L., Farnetani, M. A., Galluzzi, P., Vatti, G. P., Calabrese, F. and Morgese, G. (2003). Malformations of cortical development in neurofibromatosis type 1. *Neurology* **61**, 1799-1801.
- Birve, A., Sengupta, A. K., Beuchle, D., Larsson, J., Kennison, J. A., Rasmuson-Lestander, A. and Muller, J. (2001). Su(z)12, a novel Drosophila Polycomb group gene that is conserved in vertebrates and plants. *Development* **128**, 3371-3379.
- Burke, A. C., Nelson, C. E., Morgan, B. A. and Tabin, C. (1995). Hox genes and the evolution of vertebrate axial morphology. *Development* **121**, 333-346.
- Cao, R. and Zhang, Y. (2004). SUZ12 is required for both the histone methyltransferase activity and the silencing function of the EED-EZH2 complex. *Mol. Cell* **15**, 57-67.
- Cao, R., Wang, L., Wang, H., Xia, L., Erdjument-Bromage, H., Tempst, P., Jones, R. S. and Zhang, Y. (2002). Role of histone H3 lysine 27 methylation in Polycomb-group silencing. *Science* **298**, 1039-1043.
- Faust, C., Lawson, K. A., Schork, N. J., Thiel, B. and Magnuson, T. (1998). The Polycomb-group gene *eed* is required for normal morphogenetic movements during gastrulation in the mouse embryo. *Development* **125**, 4495-4506.

- Feinberg, A. P.** (2007). Phenotypic plasticity and the epigenetics of human disease. *Nature* **447**, 433-440.
- Kawai, J., Shinagawa, A., Shibata, K., Yoshino, M., Itoh, M., Ishii, Y., Arakawa, T., Hara, A., Fukunishi, Y., Konno, H. et al.** (2001). Functional annotation of a full-length mouse cDNA collection. *Nature* **409**, 685-690.
- Kessel, M. and Gruss, P.** (1990). Murine developmental control genes. *Science* **249**, 374-379.
- Kibar, Z., Capra, V. and Gros, P.** (2007). Toward understanding the genetic basis of neural tube defects. *Clin. Genet.* **71**, 295-310.
- Korf, B. R., Schneider, G. E. and Poussaint, T. Y.** (1999). Structural anomalies revealed by neuroimaging studies in the brains of patients with neurofibromatosis type 1 and large deletions. *Genet. Med.* **1**, 136-140.
- Lasorella, A., Stegmuller, J., Guardavaccaro, D., Liu, G., Carro, M. S., Rothschild, G., de la Torre-Ubieta, L., Pagano, M., Bonni, A. and Iavarone, A.** (2006). Degradation of Id2 by the anaphase-promoting complex couples cell cycle exit and axonal growth. *Nature* **442**, 471-474.
- Lee, T. I., Jenner, R. G., Boyer, L. A., Guenther, M. G., Levine, S., Kumar, R. M., Chevalier, B., Johnstone, S. E., Cole, M. F., Isono, K. et al.** (2006). Control of developmental regulators by Polycomb in human embryonic stem cells. *Cell* **125**, 301-313.
- Lennon, G., Auffray, C., Polymeropoulos, M. and Soares, M. B.** (1996). The I.M.A.G.E. Consortium: an integrated molecular analysis of genomes and their expression. *Genomics* **33**, 151-152.
- Nagy, A., Gersenstein, M. and Vintersten, K.** (2003). *Manipulating The Mouse Embryo: A Laboratory Manual*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- O'Carroll, D., Erhardt, S., Pagani, M., Barton, S. C., Surani, M. A. and Jenuwein, T.** (2001). The polycomb-group gene Ezh2 is required for early mouse development. *Mol. Cell. Biol.* **21**, 4330-4336.
- Parmar, H., Patkar, D., Shah, J. and Maheshwari, M.** (2003). Diastematomyelia with terminal lipomyelocystocele arising from one hemicord: case report. *Clin. Imaging* **27**, 41-43.
- Pasini, D., Bracken, A. P., Jensen, M. R., Lazzarini Denchi, E. and Helin, K.** (2004). Suz12 is essential for mouse development and for EZH2 histone methyltransferase activity. *EMBO J.* **23**, 4061-4071.
- Rodriguez-Henche, N., Jamen, F., Leroy, C., Bockaert, J. and Brabet, P.** (2002). Transcription of the mouse PAC1 receptor gene: cell-specific expression and regulation by Zac1. *Biochim. Biophys. Acta* **1576**, 157-162.
- Schuettengruber, B., Chourrout, D., Vervoort, M., Leblanc, B. and Cavalli, G.** (2007). Genome regulation by polycomb and trithorax proteins. *Cell* **128**, 735-745.
- Sparmann, A. and van Lohuizen, M.** (2006). Polycomb silencers control cell fate, development and cancer. *Nat. Rev. Cancer* **6**, 846-856.
- Stevenson, K. L.** (2004). Chiari Type II malformation: past, present, and future. *Neurosurg. Focus* **16**, E5.
- Suh, J., Lu, N., Nicot, A., Tatsuno, I. and DiCicco-Bloom, E.** (2001). PACAP is an anti-mitogenic signal in developing cerebral cortex. *Nat. Neurosci.* **4**, 123-124.
- Tan, J., Yang, X., Zhuang, L., Jiang, X., Chen, W., Lee, P. L., Karuturi, R. K., Tan, P. B., Liu, E. T. and Yu, Q.** (2007). Pharmacologic disruption of Polycomb-repressive complex 2-mediated gene repression selectively induces apoptosis in cancer cells. *Genes Dev.* **21**, 1050-1063.
- Tubbs, R. S., Rutledge, S. L., Kosentka, A., Bartolucci, A. A. and Oakes, W. J.** (2004). Chiari I malformation and neurofibromatosis type 1. *Pediatr. Neurol.* **30**, 278-280.
- Valente, T. and Auladell, C.** (2001). Expression pattern of Zac1 mouse gene, a new zinc-finger protein that regulates apoptosis and cellular cycle arrest, in both adult brain and along development. *Mech. Dev.* **108**, 207-211.
- Varrault, A., Gueydan, C., Delalbre, A., Bellmann, A., Houssami, S., Aknin, C., Severac, D., Chotard, L., Kahli, M., Le Digarcher, A. et al.** (2006). Zac1 regulates an imprinted gene network critically involved in the control of embryonic growth. *Dev. Cell* **11**, 711-722.
- Venturin, M., Guarnieri, P., Natacci, F., Stabile, M., Tenconi, R., Clementi, M., Hernandez, C., Thompson, P., Upadhyaya, M., Larizza, L. et al.** (2004). Mental retardation and cardiovascular malformations in NF1 microdeleted patients point to candidate genes in 17q11.2. *J. Med. Genet.* **41**, 35-41.