

CORRECTION


A patient-derived stem cell model of hereditary spastic paraplegia with *SPAST* mutations

Greger Abrahamsen, Yongjun Fan, Nicholas Matigian, Gautam Wali, Bernadette Bellette, Ratneswary Sutharsan, Jyothy Raju, Stephen A. Wood, David Veivers, Carolyn M. Sue and Alan Mackay-Sim

There was an error published in *Dis. Model. Mech.* **6**, 489-502.

Due to a technical error with the image analysis software, the published speed unit of peroxisome movement is incorrect. The stated peroxisome speed unit ($\mu\text{m}/\text{second}$) should be changed to $\mu\text{m}/2$ seconds in two different places in the paper: (1) twice in the Results section 'Dynamics of peroxisome movement', and (2) the x -axis of Fig. 5 (the legend remains the same). The correct text and figure is shown below. This change does not alter any conclusions of the paper, which was a comparison of cells from patients and controls.

(1) The mean peroxisome speed in patient cells was 93% slower than in control cells (control, $0.172 \pm 0.001 \mu\text{m}/2$ seconds; patient, $0.160 \pm 0.001 \mu\text{m}/2$ seconds; $t=9.19$, d.f.= 24,398, $P<0.0001$).



(2) **Fig. 5. Peroxisome speeds were slower in patient and control cells.** Frequency distributions of peroxisomes in different speed classes are shown for control cells (open bars; $n=7$ individuals, 10 cells per individual) and patient cells (filled bars; $n=6$ individuals, 10 cells per individual). Peroxisome speeds were quantified every 2 seconds for 2 minutes and grouped in speed classes expressed as the percentage of peroxisomes in each speed class as a percentage of the total number of peroxisomes for each group (control cells, $n=13,871$ peroxisomes; patient cells, 10,529 peroxisomes).

The authors apologise to the readers for any confusion that this error might have caused.